

Potential role of ultrafine particles in  
associations between airborne particle mass  
and cardiovascular health

Ralph J. Delfino, Constantinos Sioutas, and Shaista Malik  
doi:10.1289/ehp.7938 (available at <http://dx.doi.org/>)  
Online 16 March 2005



# **Potential role of ultrafine particles in associations between airborne particle mass and cardiovascular health**

Ralph J. Delfino<sup>1</sup>, Constantinos Sioutas,<sup>2</sup> Shaista Malik<sup>3</sup>

<sup>1</sup> Epidemiology Division, Department of Medicine, University of California, Irvine, CA, 92697

<sup>2</sup> Department of Civil and Environmental Engineering, University of Southern California, Los Angeles, CA 90089

<sup>3</sup> Cardiology Division, Department of Medicine, University of California, Irvine, Irvine, CA, 92697

*Address correspondence to:*

Ralph J. Delfino, M.D., Ph.D.,

Epidemiology Division, Department of Medicine,

University of California, Irvine,

224 Irvine Hall, Irvine, CA 92697-7550.

Phone: (714) 824-7401;

Fax: (714) 824-4773;

Email: [rdelfino@uci.edu](mailto:rdelfino@uci.edu)

**Running Title:** Cardiovascular Health and Ultrafine Particles

**Key words:** epidemiology, toxic air pollutants, diesel, cardiovascular diseases, particle size, oxidative stress, cytokines.

**Abbreviations:**

BP : blood pressure

CAP: concentrated ambient particles

CIMT: carotid intima-media thickness

CI: confidence interval

CHD: coronary heart disease

CHF: congestive heart failure

COPD: chronic obstructive pulmonary disease

DEP: Diesel exhaust particles

EC: elemental carbon

ETS: environmental tobacco smoke

HR: heart rate

HRV: heart rate variability

IL-1 $\beta$ : interleukin 1beta

IL-6: interleukin 6

NC<sub>0.01-0.1</sub>: number concentrations of ultrafine mode particles 0.01 to 0.1  $\mu\text{m}$  in diameter

NC<sub>0.1-1</sub>: number concentrations of accumulation mode particles 0.1 to 1.0  $\mu\text{m}$  in diameter

NF $\kappa$ B: nuclear transcription factor- $\kappa$ B

NMMAPS: National Morbidity, Mortality and Air Pollution study

Odds ratio: OR

PAH: polycyclic aromatic hydrocarbon

PM: particulate matter

PM<sub>2.5</sub>: particulate matter < 2.5  $\mu\text{m}$  in aerodynamic diameter

PM<sub>10</sub>: particulate matter < 10  $\mu\text{m}$  in aerodynamic diameter

PN: particle number

ROS: reactive oxygen species

RR: relative risk

TNF- $\alpha$ : tumor necrosis factor- $\alpha$

TSP: total suspended particulates (PM approximately < 50 µm in diameter)

UFP: ultrafine particulate matter, < 0.1 µm in aerodynamic diameter

**ACKNOWLEDGMENTS:** This work was supported by grant number ES-12243 from the National Institute of Environmental Health Sciences (NIEHS), U.S. National Institutes of Health (NIH), and its contents are solely the responsibility of the author and do not necessarily represent the official views of the NIEHS, NIH. This work was also supported by the Southern California Particle Center and Supersite (SCPCS) funded by the U.S. EPA (STAR award #R82735201). This manuscript has not been subjected to the EPA peer and policy review, and therefore, does not necessarily reflect the views of the Agencies. No official endorsement should be inferred. The authors declare they have no competing financial interests.

#### **OUTLINE OF MANUSCRIPT SECTION HEADERS:**

**ABSTRACT**

**Importance**

**Evidence of Causal Pollutant Components in Epidemiologic Time Series, Cohort and Cross-sectional Studies**

**Evidence for Pathophysiological Mechanisms and Causal Components in PM-related Cardiovascular Effects**

*Cardiac Ischemia and Related Outcomes*

*Blood Pressure (BP)*

*Autonomic Control of Cardiac Rhythm*

*Systemic Inflammation and Thrombosis*

*Summary and Biological Plausibility*

**Conclusion**

**REFERENCES**

**TABLE 1**

**Figure Legend**

**FIGURE 1**

## ABSTRACT

Numerous epidemiologic time series studies have shown generally consistent associations of cardiovascular hospital admissions and mortality with outdoor air pollution, particularly mass concentrations of particulate matter (PM)  $< 2.5$  or  $< 10$   $\mu\text{m}$  in diameter (PM<sub>2.5</sub>, PM<sub>10</sub>). Panel studies with repeated measures have supported the time series results showing associations between PM and risk of cardiac ischemia and arrhythmias, increased blood pressure, decreased heart rate variability, and increased circulating markers of inflammation and thrombosis. The causal components driving the PM associations remain to be identified. Epidemiologic data using pollutant gases and particle characteristics such as particle number concentration and elemental carbon has provided indirect evidence that products of fossil fuel combustion are important. Ultrafine particles  $< 0.1$   $\mu\text{m}$  (UFP) dominate particle number concentrations and surface area, and are therefore capable of carrying large concentrations of adsorbed or condensed toxic air pollutants. It is likely that redox active components in UFP from fossil fuel combustion reach cardiovascular target sites. High UFP exposures may lead to systemic inflammation through oxidative stress responses to reactive oxygen species, and thereby promote the progression of atherosclerosis and precipitate acute cardiovascular responses ranging from increased blood pressure to myocardial infarction. The next steps in epidemiologic research are to identify more clearly the putative PM casual components and size fractions linked to their sources. To advance this, we discuss in a companion paper (Sioutas et al. in press) the need for and methods of UFP exposure assessment.

## **Importance**

Coronary heart disease (CHD) is the leading cause of death and hospitalization among adults 65 years of age or older (Desai et al. 1999), which makes the identification of preventable causes for heart disease morbidity and mortality an important research goal. Numerous epidemiologic time series studies have shown generally consistent associations of outdoor (ambient) air pollution with cardiovascular hospital admissions (Burnett et al. 1995; 1997a; 1997b; 1999; D'Ippoliti et al. 2003; Le Tertre et al. 2002; Linn et al. 2000; Mann et al. 2002; Morris et al. 1995; Peters et al. 2001a; Poloniecki et al. 1997; Samet et al. 2000a; Schwartz and Morris 1995; Schwartz 1999; Zanobetti et al. 2000a; 2000b; 2001). Consistent associations of ambient air pollution have also been found with cardiovascular mortality (Clancy et al. 2002; Dockery et al. 1993; Goldberg et al. 2001a, 2001b; Hoek et al. 2001; Kwon et al. 2001; Laden et al. 2000; Pope et al. 2004a; Rossi et al. 1999; Samet et al. 2000b; Schwartz et al. 1996; Wichmann et al, 2000; Zanobetti et al. 2003). The National Research Council's (NRC) Committee on Research Priorities for Airborne Particulate Matter (PM) has identified research priorities to explain the morbidity and mortality associations in the time series studies (NRC, 1998; 1999; 2001; 2004). One priority is to identify the pathophysiological mechanisms and causal pollutant components driving these associations (Seaton et al. 1995).

The causal components driving the PM relationship to cardiovascular morbidity and mortality remain to be identified. Historically, the difficulty in accomplishing this in epidemiologic studies is related to the common use of ambient air pollution data from monitoring stations located at central regional sites. This has led to both exposure misclassification and high correlations between different pollutants. Both of these problems can be addressed with measurements of personal and/or microenvironmental exposures (Sarnat et al. 2000; 2001). Another problem is that the importance of particle size and chemistry has been limited by reliance on the same government

monitoring data. In the U.S., this data generally only includes particle mass concentrations in air at two particle size cuts,  $PM_{10}$  ( $PM < 10 \mu m$  in aerodynamic diameter) and more recently  $PM_{2.5}$  ( $PM < 2.5 \mu m$ ). However, there is sufficient reason to believe that ultrafine particles ( $PM < 0.1 \mu m$ ) are important in morbidity and mortality associations otherwise attributed to larger size fractions.

Major characteristics of ultrafine particles (UFP) that support their potential importance include a high pulmonary deposition efficiency, magnitudes higher particle number concentration than larger particles and thus a much higher surface area. The ultrafine particle's surface can carry large amounts of adsorbed or condensed toxic air pollutants (oxidant gases, organic compounds and transition metals) (Oberdörster 2001). Many of these toxic air pollutants have been identified as having pro-inflammatory effects in part through the action of reactive oxygen species (ROS), but relevant exposure data is rarely available to epidemiologists. Available surrogate measures of fossil fuel combustion such as elemental carbon (EC) or black smoke are of some use in this regard. Results from a study in southern California showed a large proportion of urban UFP is made up of primary combustion products from mobile source emissions (particularly diesel and automobile exhaust), and includes organic compounds, EC and metals (Kim et al. 2002). Because exposure to mobile emissions can be variable across short distances and depends on personal activity patterns, assessing such exposures requires methods that go beyond the use of government monitoring data alone. These issues regarding the characteristics of UFP are more thoroughly discussed in a companion paper (Sioutas et al. in press)

In the present review, we discuss evidence for adverse effects of air pollution on cardiovascular health with an emphasis on findings that suggest a role for UFP and related toxic air pollutant components. To date, there is little direct epidemiologic data on UFP. Studies using other particle size fractions, other particle measurements such as black smoke, and gas-phase pollutants

are used to provide a rationale for investigations of UFP. The focus of this paper is on epidemiologic studies following individual subjects over time. Several excellent reviews of experimental data and methods can be found elsewhere (Donaldson et al. 2001; Utell et al. 2002).

### **Evidence of Causal Pollutant Components in Epidemiologic Time Series, Cohort and Cross-sectional Studies**

The National Morbidity, Mortality and Air Pollution study (NMMAPS) is the largest of the air pollution time series studies to date (Samet et al. 2000a; 2000b). Results show positive associations of PM<sub>10</sub> with cardiopulmonary mortality and with hospital admissions for cardiovascular disease, chronic obstructive pulmonary disease (COPD) and pneumonia in patients 65 years of age and older living in varied environments across up 90 cities in the U.S. A subsequent analysis to correct for statistical errors showed an increase of 0.34% (95% CI: 0.1, 0.57) in combined cardiorespiratory mortality for each 10 µg/m<sup>3</sup> of air increase in PM<sub>10</sub> (Dominici et al. 2003). Another reanalysis of hospitalizations in 14 U.S. cities by Janssen et al. (Janssen et al. 2002) broke down the PM<sub>10</sub> concentrations using information on source categories. They found that for cardiovascular admissions, and to a lesser extent COPD admissions, PM<sub>10</sub> from highway vehicle and diesel emissions and from oil combustion showed the strongest associations with the most stable regression coefficients in co-regressions with other source categories. These findings are supported by an analysis of PM data collected for the Harvard Six Cities Study (Dockery et al. 1993) by Laden et al. (2000) using elemental profiles of PM<sub>2.5</sub> samples. They showed that associations between mobile source (largely traffic-related) particles and daily total mortality for the six metropolitan areas were twice that for sulfate-rich coal combustion particles. This difference was most clearly demonstrated for deaths from CHD.



Additional information regarding causal pollutant components has come from analyses of ambient gaseous air pollutants under U.S. federal regulation (CO, NO<sub>2</sub>, SO<sub>2</sub> and O<sub>3</sub>). They can be strongly correlated with PM in ambient air. A European study by Katsouyanni et al. (2001) of 29 cities showed a positive association between total mortality and PM<sub>10</sub>, and this association was not confounded by SO<sub>2</sub> or O<sub>3</sub>. However, they did find that in cities with higher versus lower average NO<sub>2</sub>, the association with PM<sub>10</sub> was significantly greater (0.80 vs. 0.19% increase in mortality per 10 µg/m<sup>3</sup> PM<sub>10</sub>, respectively). The NMMAPS study found PM<sub>10</sub> associations with mortality were largely independent of NO<sub>2</sub>, SO<sub>2</sub> and O<sub>3</sub> (Samet et al. 2000a). Goldberg et al. (2001a, 2001b), Moolgavkar et al. (2000), and Venner et al. (2003) have also found robust associations between cardiovascular mortality and pollutant gases that were often are stronger than particle associations. In a time series study of the Los Angeles air basin, Linn et al. (2000) found significant associations of daily cardiovascular hospital admissions were strongest for CO, followed by NO<sub>2</sub> then much weaker associations for PM<sub>10</sub>, but daily PM data were limited by fewer stations. Morris et al. (1995) and Morris and Naumova (1998) found hospital admissions for congestive heart failure (CHF) were associated with CO independent of other gaseous pollutants in several large U.S. cities. Mann et al. (2002) also found significant associations of daily CHD hospital admissions with NO<sub>2</sub> and CO in Los Angeles, particularly among cases with a secondary diagnosis of CHF or arrhythmia. Lin et al. (2003) found that an interquartile range increase in CO was associated with an increase of 6.4% in daily angina and acute MI emergency room visits in Sao Paulo, Brazil. A time series study of seven European areas found cardiovascular hospital admissions, especially CHD, were associated with SO<sub>2</sub> (Sunyer et al. 2003). Associations between gases and hospital admissions for CHD and CHF have been found in several other studies (e.g., Burnett 1997b; 1999; Koken et al. 2003; Morris et al. 1995; 1998).

Some of the time series investigators have hypothesized that pollutant gases could be acting as indicators for a causal mixture of pollutants including PM-related components. Ambient CO is highly correlated with UFP near combustion sources such as freeways (discussed more fully below). Although it is possible that some of the effects detected with CO are due to the formation of carboxyhemoglobin in the blood and carboxymyoglobin in muscle, reported ambient concentrations are low ( $< 6$  ppm). A postulated mechanism for increased susceptibility to low CO doses is the attainment of a nominal threshold of reduced O<sub>2</sub> transport to the heart and further compromised cardiac myoglobin, particularly in CHF patients (McGrath, 2000).

Additional evidence of causal components linked to UFP come from European studies that have used a non-gravimetric PM measure called black smoke, which is roughly representative of EC. Le Tertre et al. (2002) conducted a time series analysis of cardiovascular hospital admissions in eight European cities and found that CHD admissions were associated with PM<sub>10</sub> and black smoke. The association with PM<sub>10</sub>, but not with black smoke, was reduced by adding CO to the model and eliminated by adding NO<sub>2</sub>. Both Le Tertre et al. (2002) and the European study by Katsouyanni et al. (2001) reported above, hypothesized that their results were attributable to traffic exhaust and its consequent high emissions of CO, NO<sub>2</sub>, black smoke, and air toxics. It is relevant to point out that traffic exhaust, particularly from diesel engines, is a major contributor to ultrafine particle mass in urban areas (Kittelson, 1998; Tobias et al. 2001) and in general, UFP are both strongly linked to mobile source emissions and laden with toxic constituents (Shi et al. 2001; Kim et al. 2002).

While time series investigations have provided important information regarding the overall public health impact of ambient air pollutants on severe outcomes such as mortality, studies of individual subjects have provided insights into the underlying acute or chronic exposure-response

relationships. Below we review studies of individuals using various epidemiologic designs, including cohort and panel studies, focusing only on findings for cardiovascular outcomes. Details for selected studies are presented in Table 1 and follow the discussion in the text.

Time series studies have provided evidence for acute effects of air pollutants on cardiovascular morbidity and mortality. However, there are still gaps in the literature regarding chronic health impacts from long-term pollutant exposures. Cohort studies are best suited to address this gap. Dockery et al. (1993) reported evidence from the Harvard Six Cities study that ambient PM<sub>2.5</sub> was associated with risk of cardiopulmonary mortality in a cohort of 8,111 adults (Table 1). Pope et al (2004a) used 16 years of data from over 500,000 adults in 151 U.S. cities who participated in the American Cancer Society II cohort. The authors found that a 10 µg/m<sup>3</sup> elevation in PM<sub>2.5</sub> was associated with 8-18% increases in mortality due to ischemic heart disease, dysrhythmias, heart failure, and cardiac arrest. Mortality from various respiratory causes was not associated with PM<sub>2.5</sub> (Table 1). In contrast, in a cohort study of 6,338 Seventh Day Adventists living in California, found associations of long-term exposure to PM and O<sub>3</sub> with respiratory mortality, but not with cardiovascular mortality (Abbey et al. 1999) (Table 1). Differences in findings might be due to exposure misclassification from the use of central regional air pollutant data. Hoek et al. (2002) tried to address this issue by evaluating effects of traffic exposures near the home in a cohort study of 5,000 adults followed 8 years in the Netherlands (Table 1). They showed that living near a major road was more strongly associated with cardiopulmonary mortality than ambient background air pollutant levels. This finding suggests that pollutants more closely associated with traffic, which include UFP and associated toxic air pollutants, could be causal components in the mortality associations.

Kunzli et al (2004) conducted a cross-sectional study of 798 healthy adults with elevated LDL-cholesterol or homocysteine living in Los Angeles (Table 1). Subjects were in a dietary supplement clinical trial with ultrasound data on carotid intima-media thickness (CMT) as an estimate of atherosclerosis. Exposure included an estimate of using a geostatistical model to link subject address to annual mean PM<sub>2.5</sub> from 23 local air monitoring stations. They found positive associations between CMT and PM<sub>2.5</sub>, adjusting for host risk factors. Associations were larger for women, older subjects ( $\geq 60$ ), subjects on lipid lowering medications, and never smokers.

### **Evidence for Pathophysiological Mechanisms and Causal Components in PM-related Cardiovascular Effects**

The following section looks at epidemiologic studies called panel studies, which are designed to evaluate the relationship between repeated air pollutant exposures and cardiovascular outcomes in individual subjects. We augment this discussion with a few selected human clinical studies that extend the panel studies findings using controlled exposures, particularly those that aim to replicate ambient air mixtures. The discussion is divided by related groups of cardiovascular outcomes.

#### ***Cardiac Ischemia and Related Outcomes***

Only one published study to our knowledge has examined evidence for the relationship of particulate air pollutant exposure to cardiac ischemia in humans. An epidemiologic study of 45 adults with stable CHD conducted by Pekkanen et al. (2002) analyzed data from repeated biweekly in-clinic ECG measurements during submaximal exercise testing and outdoor ultrafine and fine particles measured at a central regional site of Helsinki, Finland (Table 1). They found significant associations between risk of ST segment depression and ambient PM<sub>2.5</sub> mass, number

concentrations of ultrafine mode particles 0.01 to 0.1  $\mu\text{m}$  in diameter ( $\text{NC}_{0.01-0.1}$ ) and number concentrations of accumulation mode particles 0.1 to 1.0  $\mu\text{m}$  in diameter ( $\text{NC}_{0.1-1}$ ) (Table 1). Odds ratios were around 3.0 for all particle metrics for an increase around their interquartile distribution. Smaller but significant associations were also found for the gases  $\text{NO}_2$  and CO, which were moderately correlated with the co-located particle measurements. The association with ultrafine particle number concentration was independent of  $\text{PM}_{2.5}$  mass concentration. It is surprising that associations for outdoor ambient  $\text{NC}_{0.01-0.1}$  were as strong as  $\text{PM}_{2.5}$  given the expectation that human exposure to UFP is less consistently represented by central site PM monitoring than  $\text{PM}_{2.5}$ , which shows much lower spatial variability than UFP (reviewed by Pekkanen and Kulmala 2004; and Sioutas et al. in press).

Cardiorespiratory symptoms potentially related to cardiac ischemia were assessed by de Hartog et al (2003) in elderly patients with CHD. The authors found that although chest pain was not associated with PM exposure, a 10  $\mu\text{g}/\text{m}^3$  increase in ambient  $\text{PM}_{2.5}$  was associated with shortness of breath and avoidance of activities (Table 1).

A case-crossover study of 691 subjects from the Augsburg Myocardial Infarction Registry found a two to three times increased risk of myocardial infarction (MI) for time-activity diary reports of hours exposed to traffic, particularly for times spent in cars and public transportation in the hours leading up to cardiac symptom onset (Peters et al. 2004) (Table 1). No direct air pollutant measurements were available. However, as discussed in our companion paper, exposures to UFP can be magnitudes higher than background levels within vehicles and near busy highways, and to a much greater degree than larger particles (Sioutas et al. in press). Accumulation mode PM, volatile organic compounds and gases such as CO could have also played a role in the findings of Peters et al. (2004).

### ***Blood Pressure (BP)***

Two studies showing associations between air pollution and BP followed subjects with COPD (Linn et al. 1999; Brauer et al. 2001, Table 1). Linn et al. (1999) found that for only 120 total person-observation times in 30 subjects, an increase of  $33 \mu\text{g}/\text{m}^3$  ambient  $\text{PM}_{10}$  (study mean) was associated with a 5.7 mm Hg increase in systolic BP. In contrast, Brauer et al. (2001) found systolic BP was inversely, but weakly associated with personal  $\text{PM}_{2.5}$  in a pooled regression analysis of 16 subjects with COPD monitored on seven separate days. This association was not confounded by inverse associations with ambient CO. Inverse associations with ambient  $\text{PM}_{10}$  were larger, but confounded by CO. Another study examined 2,607 German adults less than 65 years of age evaluated on two occasions three years apart and found a positive association of systolic BP with ambient concentrations of both total suspended particulates (TSP) and  $\text{SO}_2$  (Ibald-Mulli et al, 2001) (Table 1).

Ibald-Mulli et al. (2004) conducted one of the few panel studies to focus on the relationship between UFP and BP (Table 1). They followed 131 adults with CHD in three European centers every two weeks for around 11 clinic visits. An increase of a 5-d average of  $10,000/\text{cm}^3$  ultrafine particles ( $\text{PM}_{0.01-0.1}$ ) was associated with small decrease in systolic BP ( $-0.72 \text{ mm Hg}$ ,  $p < 0.01$ ) and diastolic BP ( $-0.70 \text{ mm Hg}$ ,  $p < 0.01$ ). Comparably small associations were also found for CO,  $1,000/\text{cm}^3$  of accumulation mode particles and  $10 \mu\text{g}/\text{m}^3$   $\text{PM}_{2.5}$ . Authors hypothesized that BP medications in these CHD patients might have blunted or modified the response to air pollution exposure. However, these results contrast those of a panel study by Zanobetti et al. (2004), who found ambient 5-d average  $\text{PM}_{2.5}$  was positively associated with blood pressure among 62 patients with pre-existing heart disease, using data from 631 repeated visits for cardiac rehabilitation in Boston (Table 1).

Panel study results for PM<sub>2.5</sub> can be compared to two experimental human studies (Brook et al., 2002; Gong et al. 2003; not shown in Table 1). Gong et al. (2003) studied the effects of PM<sub>2.5</sub> concentrated ambient particles (CAP) from Los Angeles air versus clean air on systolic BP in 12 healthy versus 12 asthmatic adults using a 2-hr rest-exercise exposure period in a chamber. CAPs are used to approximate the effects of “real world” particles. They found inverse associations of PM<sub>2.5</sub> CAP with systolic BP in asthmatics, but positive associations in healthy subjects. Results from two small studies by Brauer et al. (2001) and Gong et al. (2003) with relatively good exposure data show that PM<sub>2.5</sub> mass is inversely associated with BP in subjects with obstructive lung diseases. Brook et al., (2002) also studied the vascular effects of 150 µg/m<sup>3</sup> PM<sub>2.5</sub> CAP from Toronto air, adding 120 ppb O<sub>3</sub>, in 25 healthy adults using a 2-hr exposure period in a chamber. They found a significant but small 0.1 mm decrease in brachial artery diameter by ultrasonography for the joint exposures versus filtered air, but no change in BP, flow-mediated diameter (endothelium dependent) or nitroglycerin-mediated dilatation (endothelium independent). A follow-up analysis showed that the organic and elemental carbon fraction of PM<sub>2.5</sub> CAP were significant determinants of the effects on brachial artery diameter, which is a more sensitive biomarker of effect than BP (Urch et al. 2004).

Potential mechanisms for the observed PM-associated increases in BP have been suggested to include an increase in sympathetic tone and/or the modulation of basal systemic vascular tone due to increased concentrations of a plasma peptide known as endothelin-1 (Ibald-Mulli et al, 2001). Endothelin-1 has multiple cardiovascular actions, including vasoconstriction, leading to maintenance of basal vascular tone and BP (Haynes and Webb, 1998) and accentuating BP elevation in more severe, sodium-sensitive hypertension (Schiffrin, 2001). It is directly associated with the severity of CHF and risk of subsequent cardiac death in CHF patients (Tsutamoto et al.

1995; Galatius-Jensen et al. 1996). Endothelin-1 is produced and cleared in the lung and is generated in response to the presence of ROS (free radicals) and their metabolites (Haynes and Webb, 1998). This leaves open the possibility that pollutants could induce an excess production of endothelin-1. Supporting evidence is that urban particles have been shown to increase endothelin-1 in rats (Bouthillier et al. 1998). Effects of endothelin-1 are partly counterbalanced by vasodilatory influences of endothelial NO (Vanhoutte, 2000). Endothelial NO synthase produces NO, which traverses the extracellular space to induce smooth muscle relaxation in the vessel wall. One ROS that can be produced in the presence certain pollutant components is superoxide, which can react with NO to form the potent oxidant, peroxynitrite. Peroxynitrite is likely involved in lipid peroxidation (O'Donnell & Freeman, 2001). Therefore, an additional potential mechanism whereby pollutant components can increase BP includes superoxide-mediated inhibition of the actions of NO in inducing vasodilatation.

Despite the above data on potential biological mechanisms, epidemiologic studies reviewed have found both a decrease and increase in BP in relation to air pollutant exposures. This may be because of differences between subject populations, differences in the types of regional air pollutants, or possibly due to medications used or underlying pathology (healthy, COPD, asthma, CHD, etc). There is also a lack of data in most studies on other influences on blood pressure, namely emotional states and physical activity, which could have sustained influence on non-ambulatory BP measurements. The above factors could result in contrasting shifts in sympathetic and vagal tone in response to inhaled air pollutants, or contrasting shifts in the balance between mediators such as endothelin-1 and endothelial NO. The time course of exposure-response relationships is also ill defined, particularly periods of exposure averaging times range from minutes to days. None of the epidemiologic studies used ambulatory BP monitoring to assess acute effects



of real time changes in exposure. Ambulatory BP monitoring is more closely associated with end organ damage (heart, kidney, brain) than isolated SBP or DBP readings taken in clinic offices (Mancia and Parati 2000).

### ***Autonomic Control of Cardiac Rhythm***

Heart rate variability (HRV) is a widely used non-invasive method to investigate cardiovascular autonomic control. Reduced HRV has been shown to be a predictor of increased mortality after a myocardial infarction (Kleiger et al. 1987; La Rovere et al. 1998) and has been related especially to sudden arrhythmic death (Odemuyiwa et al. 1991, Hartikainen et al. 1996). Fourier analysis of HRV can show the magnitude of variance in the heart's rhythm across different frequency bands. Different autonomic influences on cardiovascular function (heart rate and BP) are reflected by different frequency bands. The high frequency band (0.15-0.40 Hz) has been used to estimate cardiac vagal control and is linked to respiratory influences (Task Force, 1996). Lower frequencies (0.04-0.15 Hz) are believed to represent mixed sympathetic and parasympathetic influences (Task Force, 1996). Time domain measurements are also employed (described below).

One controlled exposure study showed significant decreases in HRV in 10 healthy elderly adults for 2 h exposures to CAPS from Chapel Hill, NC (mostly mobile source), as compared with clean air, and the decrease persisted 24h later (Devlin et al. 2003). In epidemiologic studies discussed below, ambient PM has been associated with decreased heart rate variability (HRV) (Peters et al. 1999; Pope et al. 1999; Liao et al. 1999; Gold et al, 2000; Creason et al. 2001; Magari et al. 2001; 2002a, 2002b; Chan et al, 2004; Holguin et al, 2003, Pope et al, 2004b), and cardiac arrhythmia (Peters et al. 2000). Only two studies to our knowledge have investigated effects of

personal PM exposures on HRV (Chan et al. 2004; Magari et al. 2001) and one on personal CO (Tarkiainen et al. 2003).

Liao et al. (1999) showed that the largest inverse associations between non-ambulatory HRV measures and  $PM_{2.5}$  were for subjects with a history of cardiovascular conditions, although the number subjects (18) was small and the specific illnesses were not separated (not shown in Table 1). Another study of 56 elderly subjects showed inverse associations of non-ambulatory high and low frequency HRV with indoor and outdoor 24-hr gravimetric  $PM_{2.5}$  collected in a retirement home (Creason et al. 2001, not shown in Table 1). Using hourly ambient  $PM_{2.5}$  data, they briefly reported that models using prior 4-hr average  $PM_{2.5}$  and time lagged 4-hr  $PM_{2.5}$  were similar in magnitude to effects of the 24-hr  $PM_{2.5}$  averages, suggesting a mixture of short-term and cumulative effects. Holguin et al. (2003) studied 34 elderly nursing home residents living in Mexico City and showed a strong decrease in the high frequency component of HRV with high ambient  $PM_{2.5}$  exposure, and the association was stronger for indoor home  $PM_{2.5}$ . Those with hypertension had the largest reductions in HRV (Table 1). Pope et al (1999) also used ambulatory HR monitoring in seven elderly subjects with respiratory and cardiovascular disease before, during, and after episodes of elevated pollution. They found ambient  $PM_{10}$  was associated with decreased in the SD of normal RR intervals (SDNN), a time domain measure of overall HRV. However, they also found but an increase in the square root of the mean of squared differences between adjacent NN intervals (r-MSSD) (time domain measurement that corresponds to high frequency variability and parasympathetic tone). A larger study using ambulatory ECG monitors by Pope et al. (2004b) found that ambient  $PM_{2.5}$  was associated with a decrease in both SDNN and r-MSSD in 88 elderly subjects in Utah (Table 1). Magari et al. (2001) studied 40 workers occupationally exposed to welding fumes and residual oil fly ash with 24-hr monitoring using ambulatory heart rate (HR)

monitors and personal real-time PM<sub>2.5</sub> measurements from a TSI Inc. DustTrak (Shoreview, MN) (Table 1). They found significant decreases in standard deviation of average 5-min normal-to-normal intervals (SDANN) in relation to increases in prior 1-hr moving averages of PM<sub>2.5</sub>. They also found increasingly greater decreases in SDNN for higher PM<sub>2.5</sub> across longer PM<sub>2.5</sub> averaging times up to 9 hours. Magari et al. (2001) suggested inhaled particles directly affects autonomic function through a sympathetic stress response, represented by their acute response finding, and/or secondarily through airway inflammation and cytokine release into the circulation, represented by their cumulative response finding. Riediker et al. 2004 placed portable air-quality monitors in patrol cars of nine healthy male North Carolina Highway Patrol troopers who wore ambulatory ECG monitors (Table 1). In-vehicle PM<sub>2.5</sub> was positively associated with ectopic beats, heart beat cycle length, HF HRV and SDNN.

Chan et al. (2004) conducted the only study to date to assess the relationship between HRV and particle number concentrations (dominated by UFP) for particles 0.02-1.0 µm in diameter (NC<sub>0.02-1</sub>) (Table 1). They followed 9 young healthy adults (2 females) and 10 elderly male subjects with obstructive lung function impairment. This was also the first study to examine the effects of personal exposure to UFP on HRV. Subjects were monitored over only 10 daytime hours using a P-Trak Ultrafine Particle Counter (TSI Inc., Shoreview, MN) for NC<sub>0.02-1</sub>. Subjects also wore ambulatory ECG monitors for continuous 5-min beat-to-beat intervals to assess HRV. Using linear mixed effects models they found decreases in HRV indices (SDNN and r-MSSD) were associated with exposure to one to four-hr moving averages of NC<sub>0.02-1</sub> before the 5-min HRV measurements, adjusting for age, sex, body mass index, environmental tobacco smoke exposure and temperature (Table 1). Associations were stronger for the elderly panel, with the strongest effects from two-hr average NC<sub>0.02-1</sub>. These results along with Magari et al. (2001) suggest that effect of personal PM

exposure on autonomic function is acute, although the monitoring period (10 hr) was too short in Chan et al (2004) to assess longer-term effects.

Tarkiainen et al. (2003) studied six patients with CHD for one day per week for three weeks with continuous personal CO exposure monitors, ambulatory ECG monitoring for HRV and time-activity diaries and found r-MSSD increased in relation to high CO exposures (> 2.7 ppm peaks lasting 17 min, SD 8 min) (Table 1). This result contrasted results of most studies using PM exposures except Pope et al (1999). No particle data was available, but it is again important to note that outdoor CO at sites close to dense traffic is highly correlated with UFP (Zhu et al. 2002). It is conceivable that CO and/or UFP increases vagal control and induces bradyarrhythmias.

In the one study of arrhythmias and air pollution, investigators followed 100 subjects in eastern Massachusetts with implanted defibrillators (Peters et al. 2000, Table 1). They found patients with ten or more defibrillator discharge interventions for cardiac arrhythmias experienced increased arrhythmias in association with outdoor ambient NO<sub>2</sub>, CO, black carbon, but PM<sub>2.5</sub> was less strongly related. The most robust association was found for NO<sub>2</sub>, which may have been a marker for local traffic-related pollution, whereas particle mass may have been additionally influenced by other sources. Exposure was represented by only one Boston monitoring site.

### ***Systemic Inflammation and Thrombosis***

The view that air-pollution induced airway inflammation triggers systemic hypercoagulability (Seaton et al. 1995) has been supported in recent epidemiologic studies. It is relevant in this regard that compared with unaffected people, patients with CHD (Mendall et al. 1997; Lagrand et al. 1999; Stec et al. 2000; Woods et al. 2000), or a complication of CHD, CHF (Pye et al. 1990; Torre-Amione et al. 1996) have increased levels of inflammatory cytokines such as interleukin-1 $\beta$  (IL-1 $\beta$ )

and interleukin-6 (IL-6) and tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ). They also have increased levels of circulating acute phase proteins such as CRP and fibrinogen. In patients with CHD, CRP is also a strong independent predictor of future coronary events (Rifai et al. 2001). Cohort studies have shown that levels of acute phase proteins, cytokines and hemostatic factors indicative of a thrombophilic state or endothelial activation are elevated at baseline in subjects at risk for future coronary occlusion or cardiovascular mortality (Cushman et al. 1999; Danesh et al. 2000; Folsom et al. 2001; Harris et al. 1999; Haverkate et al. 1997; Jager et al. 1999; Kuller et al. 1996; Lind et al. 2001; Malik et al. 2001; Ridker et al. 2000, 2001; Ridker 2001; Thompson et al. 1995). Air pollutant exposures that lead to acute increases in already elevated levels of inflammatory and hemostatic factors may also precipitate adverse health outcomes. This is a strong possibility in patients with diagnosed or underlying CHD, a population most likely driving the time series associations. In addition, high air pollutant exposures that lead to chronic or repeated increases in systemic inflammation through oxidative stress responses to ROS may promote the progression of atherosclerosis in susceptible individuals.

Recent studies have shown acute associations between air pollutant exposures and systemic responses indicating inflammation and hypercoagulability. Seaton et al. (1999) studied 112 elderly individuals and used one day of personal PM<sub>10</sub> data per person to predict the remaining two days using ambient (city center) PM<sub>10</sub> data (Table 1). Results showed inverse associations of estimated personal PM<sub>10</sub> with albumin-adjusted hemoglobin, packed cell volume, red blood cell count, platelets and factor VII levels. They found no associations between PM<sub>10</sub> and IL-6 or white blood cell count. Only ambient PM<sub>10</sub> was positively associated with C-reactive protein (CRP) concentrations, but it was also inversely associated with fibrinogen. The authors hypothesized that particles enter lung endothelial cells or erythrocytes and subsequently influence red cell

adhesiveness leading to peripheral sequestration of red cells. Contrasting results were found by Schwartz (2001) who used health data from the Third National Health and Nutrition Examination Survey (NHANES III) in the United States (Table 1). Results showed that outdoor PM<sub>10</sub> levels on the day of subject visits or previous day was positively associated with fibrinogen levels, and counts of platelets and white blood cells. Fibrinogen increased by 13 µg/dL (95% CI: 4.6-22.1) for an interquartile range change in PM<sub>10</sub> of 26 µg/m<sup>3</sup>. PM effects were independent of gaseous pollutants. Schwartz (2001) argued that the NHANES III results were consistent with data in controlled human exposure (Ghio et al. 2000) and animal studies (Gardner et al. 2000) that showed increased plasma fibrinogen following particle exposures. Pekkanen et al. 2000 found no association between PM<sub>10</sub> and fibrinogen using cross-sectional data from another cohort study 7205 subjects in London. However, they did find associations between fibrinogen and two pollutant gases, NO<sub>2</sub> and CO, but not SO<sub>2</sub> or O<sub>3</sub>. An epidemiologic studies in Augsburg, Germany has also shown positive associations of ambient air pollution with plasma viscosity (Peters et al, 1997a) and with C-reactive protein (CRP) concentrations (Peters et al, 2001b, Table 1). Another study of people exposed to forest fire smoke showed increased circulating levels of IL-1β and IL-6 (Van Eeden et al. 2001, not shown). A panel study by Pope et al. (2004b, Table 1) with 88 elderly subjects in Utah showed a 0.81 mg/dL CRP increase in association with a 100 µg/m<sup>3</sup> increase in ambient PM<sub>2.5</sub>. There was no association with white or red blood cell counts, platelets, or whole-blood viscosity. Riediker et al, 2004 (discussed above) assessed the relationship between in-vehicle PM exposure and markers of inflammation in nine healthy male state troopers. An in-vehicle 10 µg/m<sup>3</sup> PM<sub>2.5</sub> increase was associated with decreased lymphocytes (-11%), increased red blood cell indices (1%), neutrophils (6%), C-reactive protein (32%), and von Willebrand factor (12%).

### ***Summary and Biological Plausibility***

In summary, only three studies to date have directly evaluated impacts on cardiovascular health by UFP or particle number concentration (Chan et al. 2004; Ibalid-Mulli et al. 2004; Pekkanen et al. 2002). Results of Pekkanen et al. (2002) showing ST segment depression in relation to UFP is the most compelling finding. Associations of ambient  $NC_{0.01-0.1}$  with ST segment depression were independent of ambient  $PM_{2.5}$ , but it is unclear whether the ambient exposure data represented personal UFP exposures of subjects. Other indirect evidence that components of fossil fuel combustion are important come from studies using surrogate measures of particle composition such as black smoke, proximity of homes to traffic, or source apportionment data. Epidemiologic associations for pollutant gases also appear to add to the notion that cardiovascular effects may be linked to primary products of combustion emissions that include UFP.

Because hypertension, ST segment depression and cardiac arrhythmias are well-known risk factors for cardiac morbidity and mortality, the above findings of acute associations with PM from individual-level studies are relevant to the reported findings of time series and cohort investigations of mortality and hospital morbidity. However, mixed findings for BP have not provided a coherent view of particle effects. Findings for HRV are largely consistent in finding a decrease in HRV except for the increase in r-MSSD with ambient PM among elderly subjects found by Pope et al (1999) and increased HF HRV for in-vehicle PM among healthy men found by Riediker et al. (2004). The clinical importance of HRV to cardiovascular disease is unclear though (Task Force 1996), and many technical issues regarding the influence of respiratory patterns (respiratory sinus arrhythmia) and psychosocial stress (both unmeasured in the reviewed studies) remain unresolved (Sloan et al. 1994).

The reviewed epidemiological studies on circulating biomarkers of effect show inconsistent relationships between air pollution and blood markers of inflammation and hypercoagulability, possibly because all but two studies used ambient exposure to PM. Seaton et al. (1999) and Riediker et al. (2004) are currently the only studies that used any personal PM exposure measurements, but results are not consistent. In addition, the reviewed studies of circulating biomarkers did not target people with cardiovascular diseases, who are expected to be among the most susceptible population as indicated in the time series investigations.

The main limitation of most epidemiologic studies is exposure misclassification from dependence on central site rather than personal or microenvironmental exposure data. However, studies reported above that do have personal exposure data also have limited numbers of subjects or days monitored. In general, some major methodological issues that remain involve choice of susceptible populations, personal exposure assessment, and timing of measurements to assess the temporality of exposure-dose-response relationships.

Despite the inconsistencies in epidemiologic data, there are sound postulated mechanisms that support the biological plausibility of many of the findings. Airway inflammation from PM likely involves inhalation of agents leading to the deposition or production in lung tissue of ROS. The ROS then induce subsequent oxidant injury and inflammatory responses (Pritchard et al. 1996; Schreck et al. 1991) both in the lungs and systemically. Inhalation of particle-bound airborne transition metals (copper, iron, nickel and vanadium) can lead to the production of ROS in lung tissue. Residual oil fly ash containing high concentrations of transition metals but low in organic compounds have been shown to induce *in vitro* increases in IL-6 mRNA in human epithelial cells (Quay et al. 1998). Dogs exposed to concentrated ambient particles from Boston air showed increased BAL macrophages and increased circulating neutrophils in relation to a vanadium/nickel



factor, but no associations were shown with total mass (Clarke et al. 2000). This suggests pollutant composition was important.

Organic constituents of PM are also capable of generating ROS. Nel et al. (2001) have presented evidence that polycyclic aromatic hydrocarbons (PAH) from diesel exhaust particles (DEP) and oxidized derivatives of PAHs, such as quinones, lead to the generation of ROS and subsequent oxidant injury and inflammatory responses, including the production of nuclear transcription factor- $\kappa$ B (NF $\kappa$ B). NF $\kappa$ B increases the transcription of cytokines and acute phase proteins (Schreck et al. 1991). Evidence has been presented that DEP induces a broad polyclonal activation of cytokines from an adjuvant-like activity of DEP PAH (Diaz-Sanchez 1996; 1997; Fujieda 1998; Nel et al. 1998; 2001). Human pulmonary responses to DEP include increased neutrophils and B-lymphocytes in lavage fluids, increased expression of endothelial adhesion molecules ICAM-1 and VCAM-1 in bronchial biopsies, and increased neutrophils and platelets in peripheral blood (Salvi et al. 1999). Such DEP-induced effects from oxidative stress mechanisms would be expected to lead to increased systemic hypercoagulability, but to date supporting data in humans is limited.

Epidemiologic evidence in humans that PM exposure increases biomarkers of oxidative stress in blood is limited to one study of 50 healthy young adults in Copenhagen using air samplers carried by subjects (Sorensen et al, 2003). They found a positive association between personal black carbon exposure and 2-aminoadipic semialdehyde in plasma proteins (PLAAS, a protein oxidation product). However, no association with personal PM<sub>2.5</sub> mass was found, suggesting that traffic-related causal components may have been better represented by black carbon than by particle mass. A lipid peroxidation product (malondialdehyde), as well as RBC and hemoglobin concentrations, were positively associated with PM<sub>2.5</sub> exposure in women only.

There are also plausible linkages between pulmonary and cardiovascular responses to PM. Airway inflammatory responses have been demonstrated in animals exposed to particulate air pollutants (EPA, 2003). As discussed above, there is growing evidence that airway responses may trigger systemic inflammation and hypercoagulability. In addition, PM can induce neurogenic inflammation in the lungs from activation of capsaicin-sensitive irritant receptors leading to the release of tachykinins from sensory terminals and then airway inflammation and bronchoconstriction (Veronesi and Oortgiesen, 2001). This response could then affect cardiovascular autonomic function (Carr and Udem, 2001; Yeates, 2000) but it is not yet clear to what extent these mechanisms explain epidemiologic findings of air pollutant associations with cardiac rhythm and BP. There is limited evidence for an effect of tachykinins on cardiac function (Maggi 1996). In addition, the linkage between airway inflammation, cytokine/chemokine release and autonomic stress response has not been directly demonstrated in humans. There are some *in vitro* data linking actions of proinflammatory cytokines IL-1 $\beta$  and TNF- $\alpha$  to myocardial cell changes in contractility and action potentials (DeMeules et al. 1992; Finkel et al. 1992; Li and Rozanski, 1993; Yokoyama et al. 1993) and to induction of arrhythmias (Weisensee et al. 1993).

There is experimental data indirectly supporting a linkage between cellular inflammation in the lungs and cardiovascular responses to air pollutants. An experiment in hyperlipidemic rabbits showed that intrapharyngeal installation of ambient urban PM<sub>10</sub> led an increase in circulating PMNs and caused an increase in the volume fraction of atherosclerotic lesions, which correlated with the number of alveolar macrophages that phagocytosed PM<sub>10</sub> in the lung ( $r = 0.5$ ) (Suwa et al. 2002). Particle-induced airway inflammation and translocation of ultrafine particles and other pollutants into the circulation could lead to an increase in thrombogenic and inflammatory activity in the

blood, and to a disturbance in cardiovascular function. These extra-pulmonary effects are expected to increase the risk of adverse cardiovascular outcomes such as hospitalization.

Other evidence links airway inflammation with cardiovascular effects. Cohort data has shown links of COPD with CHD risk independent of other risk factors (Jousilahti et al. 1999; Wedzicha et al. 2000), suggesting that pulmonary inflammatory processes may have pro-inflammatory effects on the vascular endothelium. This could occur in individuals with asthma or COPD who have depleted antioxidant defenses from oxidative stress as compared with normal subjects, and their defenses are further lowered during disease exacerbations (Rahman et al. 1996). Zanobetti & Schwartz (2000a) have shown that a positive association between hospital admissions for cardiovascular diseases and ambient air pollution was nearly doubled in elderly patients admitted with concurrent respiratory infections. Diabetics appear to be another susceptible group, with stronger associations between cardiovascular hospital admissions and ambient air pollution (Zanobetti & Schwartz, 2001).

Several excellent reviews of experimental data examining acute pulmonary and cardiovascular responses to inhaled ultrafine and fine particles have proposed pathophysiological mechanisms (ATS, 1999; Dhalla et al. 2000; Donaldson et al. 2001; Godleski et al. 2000; MacNee & Donaldson, 2000; Nel et al. 2001; Utell & Frampton, 2000; van Eeden and Hogg, 2002; Utell et al 2002). We have synthesized this and other data into the following proposed sequence of events for UFP that link pulmonary and cardiovascular endpoints as follows (Figure 1). Most of these mechanisms likely also apply to larger PM size fractions, particularly soluble components of PM<sub>2.5</sub>, and retained non-soluble particles in the lung that may stimulate the bone marrow to induce similar systemic responses (van Eeden and Hogg, 2002).

- 1) UFP exposure is followed by high pulmonary deposition (ICRP, 1994; Daigle et al. 2003; Chalupa et al. 2004). Ultrafine particles and associated air toxics translocate to the interstitium and gain entry into the circulation (Nemmar et al. 2002, 2004; Oberdörster et al. 2002).
- 2) Redox active components of PM lead to the production of ROS in various cells in the lungs, blood and vascular tissues.
- 3) This is followed by oxidative stress responses in pulmonary epithelium and pulmonary vascular endothelium and in extrapulmonary vascular endothelium, leading to the production of oxidized phospholipids (especially LDL), lipid peroxidation (e.g., 8-iso-PGF<sub>2</sub>α), reduced antioxidant capacity (e.g., increase in the ratio of oxidized to reduced glutathione), and the production of superoxide anions by endothelial NADPH oxidase, all of which likely contributes to atherogenesis. Genetic polymorphisms in key metabolic enzymes likely play a role in susceptibility.
- 4) Pulmonary and extrapulmonary peripheral vascular oxidative stress result in the activation and mobilization of mononuclear leukocytes and the expression of NFκB, followed by increases in pro-inflammatory cytokines (e.g., IL-1β, IL-6 and TNF-α) and endothelial cell activation.
- 5) Emigration of inflammatory cells from blood to tissue sites involves upregulation of adhesion molecules (VCAM-1, ICAM-1) on vascular endothelium and circulating leukocytes.
- 6) Increased release of cytokines by activated mononuclear cells in the lungs and in the blood leads to initiation of hepatic synthesis of acute phase proteins (e.g., CRP and fibrinogen).
- 7) A hypercoagulable state then occurs with platelet activation, hemostasis and blood clot formation followed by fibrinolytic activity; this increases the risk of a coronary event. Cytokines may also have direct effects on cardiac function.

- 8) Endothelial cell activation also leads the expression of endothelin-1, which induces vasoconstriction, and increased systolic and diastolic BP, and the expression of extracellular superoxide dismutase (SOD). SOD catalyzes superoxide ( $O_2^{\cdot-}$ ) to  $H_2O_2$ , which lowers endothelial NO-induced vasodilation. Neuroinflammatory responses involving tachykinins and catecholamines may also affect cardiovascular autonomic tone.
- 9) The systemic inflammatory response also stimulates the bone marrow to release leukocytes and platelets, and polymorphonuclear leukocytes increasingly sequester in pulmonary capillaries to induce more inflammation.

## **Conclusion**

As presented in this article, numerous studies have implicated particulate air pollution as an important contributor to morbidity and mortality from cardiovascular causes. Most of this data have been epidemiological and have utilized available air pollution data from governmental monitoring stations. Because such data are collected to meet regulatory standards, they may not meet the needs of researchers trying to understand the causal pollutant components that lead to specific adverse health effects. UFP and related toxic constituents and precursors are examples of air pollutants that have not been fully investigated, in part due to lack of available data. To date, data from epidemiologic studies indirectly implicate traffic- and other combustion-related pollutants, which include UFP. Exposure assessment issues for UFP are complex and need to be considered before undertaking epidemiologic investigations of UFP health effects (Sioutas et al. in press).

A large body of evidence shows that inflammation and oxidative stress are related to both acute changes in cardiovascular health and chronic processes including atherosclerosis. It is likely that redox active components in UFP from fossil fuel combustion reach target sites in the lungs,

vasculature, and heart to induce inflammation and oxidative stress, adding to the burden of other known lifestyle risk factors for cardiovascular disease such as diet, tobacco smoke and stress.

## REFERENCES

- Abbey DE, Nishino N, McDonnell WF, Burchette RJ, Knutsen SF, Beeson WL, et al. 1999. Long-term inhalable particles and other air pollutants related to mortality in nonsmokers. *Am J Respir Crit Care Med* 159:373-382.
- American Thoracic Society (ATS). 1999. Recommendations for Standardized Procedures for the Online and Offline Measurement of Exhaled Lower Respiratory Nitric Oxide and Nasal Nitric Oxide in Adults and Children. *Am J Respir Crit Care Med* 160:2104-2117.
- Bouthillier L, Vincent R, Goegan P, Adamson IY, Bjarnason S, Stewart M, et al. 1998. Acute effects of inhaled urban particles and ozone: lung morphology, macrophage activity, and plasma endothelin-1. *Am J Pathol* 153:1873-1884.
- Brauer M, Ebelt ST, Fisher TV, Brumm J, Petkau AJ, Vedal S. 2001. Exposure of chronic obstructive pulmonary disease patients to particles: respiratory and cardiovascular health effects. *J Expo Anal Environ Epidemiol* 11:490-500.
- Brook RD, Brook JR, Urch B, Vincent R, Rajagopalan S, Silverman F. 2002. Inhalation of fine particulate air pollution and ozone causes acute arterial vasoconstriction in healthy adults. *Circulation* 105:1534-6.
- Burnett RT, Cakmak S, Brook JR, Krewski D. 1997a. The role of particulate size and chemistry in the association between summertime ambient air pollution and hospitalization for cardiorespiratory diseases. *Environ Health Perspect* 105:614-620.
- Burnett RT, Dales R, Krewski D, Vincent R, Dann T, Brook JR. 1995. Associations between ambient particulate sulfate and admissions to Ontario hospitals for cardiac and respiratory diseases. *Am J Epidemiol* 142:15-22.

- Burnett RT, Dales RE, Brook JR, Raizenne ME, Krewski D. 1997b. Association between ambient carbon monoxide levels and hospitalizations for congestive heart failure in the elderly in 10 Canadian cities. *Epidemiol* 8:162-167.
- Burnett RT, Smith-Doiron M, Stieb D, Cakmak S, Brook JR. 1999. Effects of particulate and gaseous air pollution on cardiorespiratory hospitalizations. *Arch Environ Health* 54:130-139.
- Carr MJ, Udem BJ. 2001. Inflammation-induced plasticity of the afferent innervation of the airways. *Environ Health Perspect* 109(Suppl 4):567-571.
- Chalupa DC, Morrow PE, Oberdorster G, Utell MJ, Frampton MW. 2004. Ultrafine particle deposition in subjects with asthma. *Environ Health Perspect* 112:879-82.
- Chan C-C, Chuang KJ, Shiao GM, Lin LY. 2004. Personal exposure to submicrometer particles and heart rate variability in human subjects. *Environ Health Perspect* 112:1063-1067.
- Clancy L, Goodman P, Sinclair H, Dockery DW. 2002. Effect of air-pollution control on death rates in Dublin, Ireland: an intervention study. *Lancet* 360:1210-1214.
- Clarke RW, Coull B, Reinisch U, Catalano P, Killingsworth CR, Koutrakis P, et al. 2000. Inhaled concentrated ambient particles are associated with hematologic and bronchoalveolar lavage changes in canines. *Environ Health Perspect* 108:1179-1187.
- Creason J, Neas L, Walsh D, Williams R, Sheldon L, Liao D, et al. 2001. Particulate matter and heart rate variability among elderly retirees: the Baltimore 1998 PM study. *J Expo Anal Environ Epidemiol* 11:116-122.
- Cushman M, Lemaitre RN, Kuller LH, Psaty BM, Macy EM, Sharrett AR, et al. 1999. Fibrinolytic activation markers predict myocardial infarction in the elderly. The Cardiovascular Health Study. *Arterioscler Thromb Vasc Biol* 19:493-438.



- Daigle CC, Chalupa DC, Gibb FR, Morrow PE, Oberdorster G, Utell MJ, et al. 2003. Ultrafine particle deposition in humans during rest and exercise. *Inhal Toxicol* 15:539-52.
- Danesh J, Whincup P, Walker M, Lennon L, Thomson A, Appleby P, et al. 2000. Low grade inflammation and coronary heart disease: prospective study and updated meta-analyses. *BMJ* 321:199-204.
- de Hartog JJ, Hoek G, Peters A, Timonen KL, Ibal-Mulli A, Brunekreef B, et al. 2003. Effects of fine and ultrafine particles on cardiorespiratory symptoms in elderly subjects with coronary heart disease: the ULTRA study. *Am J Epidemiol* 157:613-623.
- DeMeules JE, Pigula FA, Mueller M, Raymond SJ, Gamelli RL. 1992. Tumor necrosis factor and cardiac function. *J Trauma* 32:686-692.
- Desai MM, Zhang P, Hennessy CH. 1999. Surveillance for morbidity and mortality among older adults--United States, 1995-1996. *Mor Mortal Wkly Rep CDC Surveill Summ* 48:7-25.
- Devlin RB, Ghio AJ, Kehrl H, Sanders G, Cascio W. 2003. Elderly humans exposed to concentrated air pollution particles have decreased heart rate variability. *Eur Respir J Suppl* 40:76s-80s.
- Dhalla NS, Temsah RM, Netticadan T. 2000. Role of oxidative stress in cardiovascular diseases. *J Hypertens* 18:655-673.
- Diaz-Sanchez D, Tsien A, Casillas A, Dotson AR, Saxon A. 1996. Enhanced nasal cytokine production in human beings after *in vivo* challenge with diesel exhaust particles. *J Allergy Clin Immunol* 98:114-123.
- Diaz-Sanchez D, Tsien A, Fleming J, Saxon A. 1997. Combined diesel exhaust particulate and ragweed allergen challenge markedly enhances human *in vivo* nasal ragweed-specific IgE and skews cytokine production to a T helper cell 2-type pattern. *J Immunol* 158:2406-2413.

- D'Ippoliti D, Forastiere F, Ancona C, Agabiti N, Fusco D, Michelozzi P, et al. 2003. Air pollution and myocardial infarction in Rome: a case-crossover analysis. *Epidemiol* 14:528-35.
- Dockery DW, Pope CA, Xu X, Spengler JD, Ware JH, Fay ME, Ferris BG, Speizer FE. 1993. An association between air pollution and mortality in six U.S. cities. *N Engl J Med* 329:1753-1759.
- Dominici F, McDermott A, Daniels M, Zeger SL, Samet JM. 2003. Revised Analysis of the National Morbidity, Mortality, and Air Pollution Study, Part II: Mortality among residents of 90 cities. In: Health Effects Institute Research: Revised analysis of time series studies of air pollution and health. Boston MA: Health Effects Institute.
- Donaldson K, Stone V, Seaton A, MacNee W. 2001. Ambient particle inhalation and the cardiovascular system: potential mechanisms. *Environ Health Perspect* 109(Suppl 4):523-527.
- Fine PM, Si S, Geller MG, Sioutas C. 2004. Inferring the sources of fine and ultrafine PM at downwind receptor areas in the Los Angeles basin using multiple continuous monitors". *Aerosol Science Technol* 38:182-195,
- Finkel MS, Oddis CV, Jacob TD, Watkins SC, Hattler BG, Simmons RL. 1992. Negative inotropic effects of cytokines on the heart mediated by nitric oxide. *Science* 257:387-9.
- Folsom AR, Aleksic N, Park E, Salomaa V, Juneja H, Wu KK. 2001. Prospective study of fibrinolytic factors and incident coronary heart disease: the Atherosclerosis Risk in Communities (ARIC) Study. *Arterioscler Thromb Vasc Biol* 21:611-617.
- Fujieda S, Diaz-Sanchez D, Saxon A. 1998. Combined nasal challenge with diesel exhaust particles and allergen induces in vivo IgE isotype switching. *Am J Respir Cell Mol Biol* 19:507-512
- Galatius-Jensen S, Wroblewski H, Emmeluth C, Bie P, Haunso S, Kastrup J. 1996. Plasma endothelin in congestive heart failure: a predictor of cardiac death? *J Card Fail* 2:71-6.

- Gardner SY, Lehmann JR, Costa DL. 2000. Oil fly ash-induced elevation of plasma fibrinogen levels in rats. *Toxicol Sci* 56:175-80.
- Ghio AJ, Kim C, Devlin RB. 2000. Concentrated ambient air particles induce mild pulmonary inflammation in healthy human volunteers. *Am J Respir Crit Care Med* 162:981-988.
- Godleski JJ, Verrier RL, Koutrakis P, Catalano P, Coull B, Reinisch U, Lovett EG, Lawrence J, Murthy GG, Wolfson JM, Clarke RW, Nearing BD, Killingsworth C. 2000. Mechanisms of morbidity and mortality from exposure to ambient air particles. Research Report / Health Effects Institute 91:5-88.
- Gold DR, Litonjua A, Schwartz J, Lovett E, Larson A, Nearing B, et al. 2000. Ambient pollution and heart rate variability. *Circulation* 101:1267-1273.
- Goldberg MS, Burnett RT, Bailer JC 3rd, Tamblyn R, Ernst P, Flegel K, et al. 2001. Identification of persons with cardiorespiratory conditions who are at risk of dying from the acute effects of ambient air particles. *Environ Health Perspect* 109(Suppl 4):487-494.
- Goldberg MS, Burnett RT, Brook J, Bailer JC 3rd, Valois MF, Vincent R. 2001. Associations between daily cause-specific mortality and concentrations of ground-level ozone in Montreal, Quebec. *Am J Epidemiol* 154:817-826.
- Gong H Jr, Linn WS, Sioutas C, Terrell SL, Clark KW, Anderson KR, et al. 2003. Controlled exposures of healthy and asthmatic volunteers to concentrated ambient fine particles in Los Angeles. *Inhal Toxicol* 15:305-325.
- Harris TB, Ferrucci L, Tracy RP, Corti MC, Wacholder S, Ettinger WH Jr, et al. 1999. Associations of elevated interleukin-6 and C-reactive protein levels with mortality in the elderly. *Am J Med* 106:506-512.

- Hartikainen JEK, Malik M, Stauton A, Poloniecki J, Camm J. 1996. Distinction between arrhythmic and nonarrhythmic death after acute myocardial infarction based on heart rate variability, signal-averaged electrocardiogram, ventricular arrhythmias and left ventricular ejection fraction. *J Am Coll Cardiol* 28:296-304.
- Haverkate F, Thompson SG, Pyke SD, Gallimore JR, Pepys MB. 1997. Production of C-reactive protein and risk of coronary events in stable and unstable angina. European Concerted Action on Thrombosis and Disabilities Angina Pectoris Study Group. *Lancet* 349:462-466.
- Haynes WG, Webb DJ. 1998. Endothelin as a regulator of cardiovascular function in health and disease. *J Hypertens* 16:1081-1098.
- Hoek G, Brunekreef B, Fischer P, van Wijnen J. 2001. The association between air pollution and heart failure, arrhythmia, embolism, thrombosis, and other cardiovascular causes of death in a time series study. *Epidemiol* 12:355-357.
- Hoek G, Brunekreef B, Goldbohm S, Fischer P, van den Brandt PA. 2002. Association between mortality and indicators of traffic-related air pollution in the Netherlands: a cohort study. *Lancet* 360:1203-1209.
- Holguin F, Tellez-Rojo MM, Hernandez M, Cortez M, Chow JC, Watson JG, et al. 2003. Air pollution and heart rate variability among the elderly in Mexico City. *Epidemiol* 14:521-527.
- Ibald-Mulli A, Stieber J, Wichmann HE, Koenig W, Peters A. 2001. Effects of air pollution on blood pressure: a population-based approach. *Am J Public Health* 91:571-577.
- Ibald-Mulli A, Timonen KL, Peters A, Heinrich J, Wolke G, Lanki T, et al. 2004. Effects of particulate air pollution on blood pressure and heart rate in subjects with cardiovascular disease: a multicenter approach. *Environ Health Perspect* 112:369-377.

- International Commission on Radiological Protection (ICRP). 1994. Human Respiratory Tract Model for Radiological Protection. Tarrytown, NY:Elsevier Science Ltd.
- Jager A, van Hinsbergh VW, Kostense PJ, Emeis JJ, Yudkin JS, Nijpels G, et al. 1999. von Willebrand factor, C-reactive protein, and 5-year mortality in diabetic and nondiabetic subjects: the Hoorn Study. *Arterioscler Thromb Vasc Biol*; 19:3071-3078.
- Janssen NA, Schwartz J, Zanobetti A, Suh HH. 2002. Air conditioning and source-specific particles as modifiers of the effect of PM<sub>10</sub> on hospital admissions for heart and lung disease. *Environ Health Perspect* 110:43-49.
- Jousilahti P, Salomaa V, Rasi V, Vahtera E. 1999. Symptoms of chronic bronchitis, haemostatic factors, and coronary heart disease risk. *Atherosclerosis* 142:403-407.
- Katsouyanni K, Touloumi G, Samoli E, Gryparis A, Le Tertre A, Monopolis Y, et al. 2001. Confounding and effect modification in the short-term effects of ambient particles on total mortality: results from 29 European cities within the APHEA2 project. *Epidemiol* 12:521-531.
- Kim S, Shen S, Sioutas C. 2002. Size distribution and diurnal and seasonal trends of ultrafine particles in source and receptor sites of the Los Angeles basin. *J Air Waste Manage Assoc* 52:297-307.
- Kittelson DB. 1998. Engines and nanoparticles: a review. *J Aerosol Sci* 29:575-588.
- Kleiger RE, Miller JP, Bigger JT Jr, Moss AJ. 1987. Decreased heart rate variability and its association with increased mortality after acute myocardial infarction. *Am J Cardiol* 59:256-262.

- Koken PJ, Piver WT, Ye F, Elixhauser A, Olsen LM, et al. 2003. Temperature, air pollution, and hospitalization for cardiovascular diseases among elderly people in Denver. *Environ Health Perspect* 111:1312-7.
- Kuller LH, Tracy RP, Shaten J, Meilahn EN. 1996. Relation of C-reactive protein and coronary heart disease in the MRFIT nested case-control study. Multiple Risk Factor Intervention Trial. *Am J Epidemiol* 144:537-547.
- Künzli N, Jerrett M, Mack WJ, Beckerman B, LaBree L, Gilliland F, et al. 2004. Ambient air pollution and atherosclerosis in Los Angeles. *Environ Health Perspect* doi:10.1289/ehp.7523. [Online 22 November 2004]
- Kwon HJ, Cho SH, Nyberg F, Pershagen G. 2001. Effects of ambient air pollution on daily mortality in a cohort of patients with congestive heart failure. *Epidemiol* 12:413-419.
- Kwon HJ, Cho SH, Nyberg F, Pershagen G.. Effects of ambient air pollution on daily mortality in a cohort of patients with congestive heart failure. *Epidemiology* 2001 Jul;12(4):413-9.
- La Rovere MT, Bigger JT Jr, Marcus FI, Mortara A, Schwartz PJ. 1998. Baroreflex sensitivity and heart-rate variability in prediction of total cardiac mortality after myocardial infarction. ATRAMI (Autonomic Tone and Reflexes After Myocardial Infarction) Investigators. *Lancet* 351:478-484.
- Laden F, Neas LM, Dockery DW, Schwartz J. 2000. Association of fine particulate matter from different sources with daily mortality in six U.S. cities. *Environ Health Perspect* 108:941-947.
- Lagrand WK, Visser CA, Hermens WT, Niessen HW, Verheugt FW, Wolbink GJ, et al. 1999. C-reactive protein as a cardiovascular risk factor: more than an epiphenomenon? *Circulation* 100:96-102.

- Le Tertre A, Medina S, Samoli E, Forsberg B, Michelozzi P, Boumghar A, et al. 2002. Short-term effects of particulate air pollution on cardiovascular diseases in eight European cities. *J Epidemiol Community Health* 56:773-779.
- Li YH, Rozanski GJ. 1993. Effects of human recombinant interleukin-1 on electrical properties of guinea pig ventricular cells. *Cardiovasc Res* 27:525-530.
- Liao D, Creason J, Shy C, Williams R, Watts R, Zweidinger R. 1999. Daily variation of particulate air pollution and poor cardiac autonomic control in the elderly. *Environ Health Perspect* 107:521-525.
- Lin CA, Amador Pereira LA, de Souza Conceicao GM, Kishi HS, Milani R Jr, Ferreira Braga AL, et al. 2003. Association between air pollution and ischemic cardiovascular emergency room visits. *Environ Res* 92:57-63.
- Lind P, Hedblad B, Stavenow L, Janzon L, Eriksson KF, Lindgarde F. 2001. Influence of plasma fibrinogen levels on the incidence of myocardial infarction and death is modified by other inflammation-sensitive proteins: a long-term cohort study. *Arterioscler Thromb Vasc Biol* 21:452-458.
- Linn WS, Gong H Jr, Clark KW, Anderson KR. 1999. Day-to-day particulate exposures and health changes in Los Angeles area residents with severe lung disease. *J Air Waste Manag Assoc* 49(9 Spec No):108-115.
- Linn WS, Szlachcic Y, Gong H Jr, Kinney PL, Berhane KT. 2000. Air pollution and daily hospital admissions in metropolitan Los Angeles. *Environ Health Perspect* 108:427-434.
- MacNee W, Donaldson K. 2000. How can ultrafine particles be responsible for increased mortality? *Monaldi Arch Chest Dis* 55:135-139.

- Magari SR, Hauser R, Schwartz J, Williams PL, Smith TJ, Christiani DC. 2001. Association of heart rate variability with occupational and environmental exposure to particulate air pollution. *Circulation* 104:986-991.
- Magari SR, Schwartz J, Williams PL, Hauser R, Smith TJ, Christiani DC. 2002a. The association between personal measurements of environmental exposure to particulates and heart rate variability. *Epidemiology* 13:305-10.
- \_\_\_\_\_. 2002b. The association of particulate air metal concentrations with heart rate variability. *Environ Health Perspect* 110:875-880.
- Maggi CA. 1996. Tachykinins in the autonomic nervous system. *Pharmacol Res* 33:161-170.
- Malik I, Danesh J, Whincup P, Bhatia V, Papacosta O, Walker M, et al. 2001. Soluble adhesion molecules and prediction of coronary heart disease: a prospective study and meta-analysis. *Lancet* 358:971-976.
- Mancia G, Parati G. 2000. Ambulatory blood pressure monitoring and organ damage. *Hypertension* 36:894-900.
- Mann JK, Tager IB, Lurmann F, Segal M, Quesenberry CP Jr, Lugg MM, et al. 2002. Air pollution and hospital admissions for ischemic heart disease in persons with congestive heart failure or arrhythmia. *Environ Health Perspect* 110:1247-1252.
- McGrath JJ. 2000. Biological plausibility for carbon monoxide as a copollutant in PM epidemiologic studies. *Inhal Toxicol* 12(Suppl 4):91-107.
- Mendall MA, Patel P, Asante M, Ballam L, Morris J, Strachan DP, et al. 1997. Relation of serum cytokine concentrations to cardiovascular risk factors and coronary heart disease. *Heart* 78:273-277.



- Moolgavkar SH. 2000. Air pollution and daily mortality in three U.S. counties. *Environ Health Perspect* 108:777-784.
- Morawska L, Bofinger ND, Kocis L, Nwankwoala A. 1998. Submicrometer and super micrometer particles from diesel vehicle emissions. *Environ Science Technol* 32:2033-2042.
- Morris RD, Naumova EN. 1998. Carbon monoxide and hospital admissions for congestive heart failure: evidence of an increased effect at low temperatures. *Environ Health Perspect* 106:649-53.
- Morris RD, Naumova EN, Munasinghe RL. 1995. Ambient air pollution and hospitalization for congestive heart failure among elderly people in seven large US cities. *Am J Public Health* 85:1361-1365.
- National Research Council (NRC), Committee on Research Priorities for Airborne Particulate Matter. 1998. *Research Priorities for Airborne Particulate Matter, I: Immediate Priorities and Long-Range Research Portfolio*. Washington DC: National Academy Press.
- \_\_\_\_\_. 1999. *Research Priorities for Airborne Particulate Matter, II: Evaluating Research Progress and Evaluating the Portfolio*. Washington DC: National Academy Press.
- \_\_\_\_\_. 2001. *Research Priorities for Airborne Particulate Matter, III: Early Research Progress*. Washington DC: National Academy Press.
- \_\_\_\_\_. 2004. *Research Priorities for Airborne Particulate Matter: IV. Continuing Research Progress*. Washington DC: National Academy Press.
- Nel AE, Diaz-Sanchez D, Li N. 2001. The role of particulate pollutants in pulmonary inflammation and asthma: evidence for the involvement of organic chemicals and oxidative stress. *Curr Opin Pulm Med* 7:20-26.

- Nel AE, Diaz-Sanchez D, Ng D, Hiura T, Saxon A. 1998. Enhancement of allergic inflammation by the interaction between diesel exhaust particles and the immune system. *J Allergy Clin Immunol* 102:539-54.
- Nemmar A, Hoet PH, Vanquickenborne B, Dinsdale D, Thomeer M, Hoylaerts MF, et al. 2002. Passage of inhaled particles into the blood circulation in humans. *Circulation* 105:411-414.
- Nemmar A, Hoylaerts MF, Hoet PH, Nemery B. 2004. Possible mechanisms of the cardiovascular effects of inhaled particles: systemic translocation and prothrombotic effects. *Toxicol Lett* 149:243-253.
- Oberdörster G. 2001. Pulmonary effects of inhaled ultrafine particles. *Int Arch Occup Environ Health* 74:1-8.
- Oberdörster G, Gelein RM, Ferin J, Weiss B. 1995. Association of particulate air pollution and acute mortality: involvement of ultrafine particles? *Inhal Toxicol* 7:111-124.
- Oberdörster G, Sharp Z, Atudorei V, Elder A, Gelein R, Lunts A, et al. 2002. Extrapulmonary translocation of ultrafine carbon particles following whole-body inhalation exposure of rats. *J Toxicol Environ Health A* 65:1531-1543.
- Odenmuyiwa O, Malik M, Farrell T, Bashir Y, Poloniecki J, Camm J. 1991. Comparison of the predictive characteristics of heart rate variability index and left ventricular ejection fraction for all-cause mortality, arrhythmic events and sudden death after acute myocardial infarction. *Am J Cardiol* 68:434-439.
- O'Donnell VB, Freeman BA. 2001. Interactions between nitric oxide and lipid oxidation pathways: implications for vascular disease. *Circ Res* 88:12-21.
- Pekkanen J, Kulmala M. 2004. Exposure assessment of ultrafine particles in epidemiologic time-series studies. *Scand J Work Environ Health*. 30 (Suppl 2):9-18.

- Pekkanen J, Peters A, Hoek G, Tiittanen P, Brunekreef B, de Hartog J, et al. 2002. Particulate air pollution and risk of ST-segment depression during repeated submaximal exercise tests among subjects with coronary heart disease: the Exposure and Risk Assessment for Fine and Ultrafine Particles in Ambient Air (ULTRA) study. *Circulation* 106:933-938.
- Peters A, Dockery DW, Muller JE, Mittleman MA. 2001a. Increased particulate air pollution and the triggering of myocardial infarction. *Circulation* 103:2810-2815.
- Peters A, Döring A, Wichmann HE, Koenig W. 1997a. Increased plasma viscosity during an air pollution episode: a link to mortality? *Lancet* 349:1582-1587.
- Peters A, Frohlich M, Döring A, Immervoll T, Wichmann HE, Hutchinson WL, et al. 2001b. Particulate air pollution is associated with an acute phase response in men; results from the MONICA-Augsburg Study. *Eur Heart J* 22:1198-1204.
- Peters A, Liu E, Verrier RL, Schwartz J, Gold DR, Mittleman M, et al. 2000. Air pollution and incidence of cardiac arrhythmia. *Epidemiol* 11:11-17.
- Peters A, Perz S, Döring A, Stieber J, Koenig W, Wichmann HE. 1999. Increases in heart rate during an air pollution episode. *Am J Epidemiol* 150:1094-1098.
- Peters A, von Klot S, Heier M, Trentinaglia I, Hormann A, Wichmann HE, et al. 2004. Exposure to traffic and the onset of myocardial infarction. *N Engl J Med* 351:1721-30.
- Peters A, Wichmann HE, Tuch T, Heinrich J, Heyder J. 1997b. Respiratory effects are associated with the number of ultrafine particles. *Am J Respir Crit Care Med* 155:1376-1383.
- Poloniecki JD, Atkinson RW, de Leon AP, Anderson HR. 1997. Daily time series for cardiovascular hospital admissions and previous day's air pollution in London, UK. *Occup Environ Med* 54:535-540.

- Pope CA 3rd, Burnett RT, Thurston GD, Thun MJ, Calle EE, Krewski D, et al. 2004a. Cardiovascular mortality and long-term exposure to particulate air pollution: epidemiological evidence of general pathophysiological pathways of disease. *Circulation* 109:71-77.
- Pope CA 3rd, Hansen ML, Long RW, Nielsen KR, Eatough NL, Wilson WE, et al. 2004b. Ambient particulate air pollution, heart rate variability, and blood markers of inflammation in a panel of elderly subjects. *Environ Health Perspect* 112:339-345.
- Pope CA 3<sup>rd</sup>, Verrier RL, Lovett EG, Larson AC, Raizenne ME, Kanner RE, et al. 1999. Heart rate variability associated with particulate air pollution. *Am Heart J*, 138:890-899.
- Pritchard RJ, Ghio AJ, Lehmann JR, Winsett DW, Tepper JS, Park P, et al. 1996. Oxidant generation and lung injury after particulate air pollutant exposure increase with the concentrations of associated metals. *Inhal Toxicol* 8:457-477.
- Pye M, Rae AP, Cobbe SM. 1990. Study of serum C-reactive protein concentration in cardiac failure. *Br Heart J* 63:228-230.
- Quay JL, Reed W, Samet J, Devlin RB. 1998. Air pollution particles induce IL-6 gene expression in human airway epithelial cells via NF-kappaB activation. *Am J Respir Cell Mol Biol* 19:98-106.
- Rahman I, Morrison D, Donaldson K, MacNee W. 1996. Systemic oxidative stress in asthma, COPD, and smokers. *Am J Respir Crit Care Med* 154:1055-1060.
- Ridker PM. 1999. Inflammation, atherosclerosis, and cardiovascular risk: an epidemiologic view. *Blood Coagul Fibrinolysis*. 10(Suppl 1):S9-S12.
- Ridker PM. 2001. Role of inflammatory biomarkers in prediction of coronary heart disease. *Lancet* 358:946-948.

- Ridker PM, Hennekens CH, Buring JE, Rifai N. 2000. C-reactive protein and other markers of inflammation in the prediction of cardiovascular disease in women. *N Engl J Med* 342:836-843.
- Ridker PM, Rifai N, Lowenthal SP. 2001. Rapid reduction in C-Reactive protein with cerivastatin among 785 patients with hypercholesterolemia. *Circulation* 103:1191-1193.
- Riediker M, Cascio WE, Griggs TR, Herbst MC, Bromberg PA, Neas L, et al. 2004. Particulate matter exposure in cars is associated with cardiovascular effects in healthy young men. *Am J Respir Crit Care Med* 169:934-940.
- Rifai N, Ridker PM. 2001. High-sensitivity C-reactive protein: a novel and promising marker of coronary heart disease. *Clin Chem* 47:403-411.
- Rossi G, Vigotti MA, Zanobetti A, Repetto F, Gianelle V, Schwartz J. 1999. Air pollution and cause-specific mortality in Milan, Italy, 1980-1989. *Arch Environ Health* 54:158-164.
- Salvi S, Blomberg A, Rudell B, Kelly F, Sandström T, Holgate ST, et al. 1999. Acute inflammatory responses in the airways and peripheral blood after short-term exposure to diesel exhaust in healthy human volunteers. *Am J Respir Crit Care Med* 159:702-709.
- Samet JM, Dominici F, Curriero FC, Coursac I, Zeger SL. 2000a. Fine Particulate Air Pollution and Mortality in 20 U.S. Cities, 1987-1994. *N Engl J Med* 343:1742-1749.
- Samet JM, Zeger SL, Dominici F, Curriero F, Coursac I, Dockery DW, et al. 2000b. The National Morbidity, Mortality, and Air Pollution Study. Part II: Morbidity and mortality from air pollution in the United States. *Health Effects Institute Research Report* 94(Pt 2):5-79.
- Sarnat JA, Koutrakis P, Suh HH. 2000. Assessing the relationship between personal particulate and gaseous exposures of senior citizens living in Baltimore, MD. *J Air Waste Manage Assoc* 50:1184-1198.

- Sarnat JA, Schwartz J, Catalano PJ, Suh HH. 2001. Gaseous Pollutants in Particulate Matter Epidemiology: Confounders or Surrogates? *Environ Health Perspect* 109:1053-1061.
- Schiffrrin EL. 2001. A critical review of the role of endothelial factors in the pathogenesis of hypertension. *J Cardiovasc Pharmacol* 38(Suppl 2):S3-S6.
- Schreck R, Rieber P, Baeuerle PA. 1991. Reactive oxygen intermediates as apparently widely used messengers in the activation of the NF-kappa B transcription factor and HIV-1. *EMBO J* 10:2247-2258.
- Schwartz J. 1999. Air pollution and hospital admissions for heart disease in eight U.S. counties. *Epidemiol* 10:17-22.
- Schwartz J. 2001. Air pollution and blood markers of cardiovascular risk. *Environ Health Perspect* 109(suppl 3):405-409.
- Schwartz J, Dockery DW, Neas LM. 1996. Is daily mortality associated specifically with fine particles? *J Air Waste Manag Assoc* 46:927-939.
- Schwartz J, Morris R. 1995. Air pollution and hospital admissions for cardiovascular disease in Detroit, Michigan. *Am J Epidemiol* 142:23-35.
- Seaton A, MacNee W, Donaldson K, Godden D. 1995. Particulate air pollution and acute health effects. *Lancet* 345:176-178.
- Seaton A, Soutar A, Crawford V, Elton R, McNerlan S, Cherrie J, et al. 1999. Particulate air pollution and the blood. *Thorax* 54:1027-1032.
- Shi JP, Evans DE, Khan AA, Harrison RM. 2001. Source and concentration of nanoparticles (< 10 nm diameter) in the urban atmosphere. *Atmos Environ* 35:1193-1202.
- Sioutas C, Delfino RJ, M Singh. In press. Exposure assessment for atmospheric ultrafine particles (UFP) and implications in epidemiological research. *Environ Health Perspect*.

- Sloan RP, Shapiro PA, Bagiella E, Boni SM, Paik M, Bigger JT Jr, et al. 1994. Effect of mental stress throughout the day on cardiac autonomic control. *Biol Psych* 37:89-99.
- Sorensen M, Daneshvar B, Hansen M, Dragsted LO, Hertel O, Knudsen L, et al. 2003. Personal PM<sub>2.5</sub> exposure and markers of oxidative stress in blood. *Environ Health Perspect* 111:161-165.
- Stec JJ, Silbershatz H, Tofler GH, Matheney TH, Sutherland P, Lipinska I, et al. 2000. Association of fibrinogen with cardiovascular risk factors and cardiovascular disease in the Framingham Offspring Population. *Circulation* 102:1634-1638.
- Sunyer J, Ballester F, Tertre AL, Atkinson R, Ayres JG, Forastiere F, et al. 2003. The association of daily sulfur dioxide air pollution levels with hospital admissions for cardiovascular diseases in Europe (The Aphea-II study). *Eur Heart J* 24:752-60.
- Suwa T, Hogg JC, Quinlan KB, Ohgami A, Vincent R, van Eeden SF. 2002. Particulate air pollution induces progression of atherosclerosis. *J Am Coll Cardiol* 39:935-942.
- Tarkiainen TH, Timonen KL, Vanninen EJ, Alm S, Hartikainen JEK, Pekkanen J. 2003. Effect of acute carbon monoxide exposure on heart rate variability in patients with coronary artery disease. *Clin Physiol Funct Imaging* 23:98-102.
- Task Force of the European Society of Cardiology and the North American Society of Pacing and Electrophysiology. 1996. Heart rate variability: Standards of measurement, physiological interpretation, and clinical use. *Circulation* 93:1043-1065
- Thompson SG, Kienast J, Pyke SD, Haverkate F, van de Loo JC. 1995. Hemostatic factors and the risk of myocardial infarction or sudden death in patients with angina pectoris. European Concerted Action on Thrombosis and Disabilities Angina Pectoris Study Group. *N Engl J Med* 332:635-641.

- Tobias HJ, Beving DE, Ziemann PJ, Sakurai H, Zuk M, McMurry PH, et al. 2001. Chemical analysis of diesel engine nanoparticles using a nano-DMA/thermal desorption particle beam mass spectrometer. *Environ Science Technol* 35: 2233-2243.
- Torre-Amione G, Kapadia S, Benedict C, Oral H, Young JB, Mann DL. 1996. Proinflammatory cytokine levels in patients with depressed left ventricular ejection fraction: a report from the Studies of Left Ventricular Dysfunction (SOLVD). *J Am Coll Cardiol* 27:1201-1206.
- Tsutamoto T, Hisanaga T, Fukai D, Wada A, Maeda Y, Maeda K, et al. 1995. Prognostic value of plasma soluble intercellular adhesion molecule-1 and endothelin-1 concentration in patients with chronic congestive heart failure. *Am J Cardiol* 76:803-808.
- Urch B, Brook JR, Wasserstein D, Brook RD, Rajagopalan S, Corey P, et al. 2004. Relative contributions of PM<sub>2.5</sub> chemical constituents to acute arterial vasoconstriction in humans. *Inhal Toxicol* 16:345-52.
- U.S. EPA. 2003. Air Quality Criteria for Particulate Matter, Fourth External Review Draft. EPA/600/P-99/002aD and bD. Research Triangle Park, NC: Office of Research and Development, National Center for Environmental Assessment.
- Utell MJ, Frampton MW. 2000. Acute health effects of ambient air pollution: the ultrafine particle hypothesis. *J Aerosol Med* 13:355-359.
- Utell MJ, Frampton MW, Zareba W, Devlin RB, Cascio WE. 2002. Cardiovascular effects associated with air pollution: potential mechanisms and methods of testing. *Inhal Toxicol* 14:1231-47.
- van Eeden SF, Hogg JC. 2002. Systemic inflammatory response induced by particulate matter air pollution: the importance of bone-marrow stimulation. *J Toxicol Environ Health A* 65:1597-1613.



- van Eeden SF, Tan WC, Suwa T, Mukae H, Terashima T, Fujii T, et al. 2001. Cytokines involved in the systemic inflammatory response induced by exposure to particulate matter air pollutants PM<sub>10</sub>. *Am J Respir Crit Care Med* 164:826-30.
- Vanhoutte PM. Say NO to ET. 2000. *J Auton Nerv Syst* 81:271-277.
- Venners SA, Wang B, Xu Z, Schlatter Y, Wang L, Xu X. 2003. Particulate matter, sulfur dioxide, and daily mortality in Chongqing, China. *Environ Health Perspect* 111:562-567.
- Veronesi B, Oortgiesen M. 2001. Neurogenic inflammation and particulate matter (PM) air pollutants. *Neurotoxicology* 22:795-810.
- Wedzicha JA, Seemungal TA, MacCallum PK, Paul EA, Donaldson GC, Bhowmik A, et al. 2000. Acute exacerbations of chronic obstructive pulmonary disease are accompanied by elevations of plasma fibrinogen and serum IL-6 levels. *Thromb Haemost* 84:210-215.
- Weisensee D, Bereiter-Hahn J, Schoeppe W, Low-Friedrich I. 1993. Effects of cytokines on the contractility of cultured cardiac myocytes. *Int J Immunopharmacol* 15:581-587.
- Wichmann HE, Spix C, Tuch T, Wolke G, Peters A, Heinrich J, et al. 2000. Daily mortality and fine and ultrafine particles in Erfurt, Germany part I: role of particle number and particle mass. *Res Rep Health Eff Inst* 98:5-86.
- Woods A, Brull DJ, Humphries SE, Montgomery HE. 2000. Genetics of inflammation and risk of coronary artery disease: the central role of interleukin-6. *Eur Heart J* 21:1574-1583.
- Yeates DB. 2000. Neurally mediated cardiopulmonary and systemic responses to inhaled irritants and antigens. In: *Particle-lung Interactions* (Gehr P, Heyder J, eds). New York, NY:Marcel Dekker,

- Yokoyama T, Vaca L, Rossen RD, Durante W, Hazarika P, Mann DL. 1993. Cellular basis for the negative inotropic effects of tumor necrosis factor-alpha in the adult mammalian heart. *J Clin Invest*; 92:2303-2312.
- Zanobetti A, Canner MJ, Stone PH, Schwartz J, Sher D, Eagan-Bengston E, et al. 2004. Ambient pollution and blood pressure in cardiac rehabilitation patients. *Circulation* 110:2184-9.
- Zanobetti A, Schwartz J. 2001. Are diabetics more susceptible to the health effects of airborne particles? *Am J Respir Crit Care Med* 164:831-833.
- Zanobetti A, Schwartz J, Dockery DW. 2000b. Airborne particles are a risk factor for hospital admissions for heart and lung disease. *Environ Health Perspect* 108:1071-1077.
- Zanobetti A, Schwartz J, Gold D. 2000a. Are there sensitive subgroups for the effects of airborne particles? *Environ Health Perspect*; 108:841-845.
- Zanobetti A, Schwartz J, Samoli E, Gryparis A, Touloumi G, Peacock J, et al. 2003. The temporal pattern of respiratory and heart disease mortality in response to air pollution. *Environ Health Perspect* 111:1188-93.
- Zhu Y, Hinds WC, Kim S, Sioutas C. 2002. Concentration and size distribution of ultrafine particles near a major highway. *J Air Waste Manage Assoc* 52:1032-1042.

Table 1: Cardiovascular effects<sup>a</sup> associated with personal and ambient air pollution exposure: Selected studies

Studies	Design and Population	Outcomes	Findings for PM Mass and Components	Findings for Gases
<b>Cohort and Cross-sectional Studies</b>				
Dockery et al. 1993	Cohort study examining ambient air pollution exposure and mortality in 8111 adults in six US cities with 14 to 16 years of follow-up.	Cardiopulmonary mortality	Compared to the least polluted city, the most polluted city had an adjusted RR for cardiopulmonary mortality of 1.37 (95% CI: 1.11, 1.68).	No association with O <sub>3</sub> , but SO <sub>2</sub> and NO <sub>2</sub> tracked between-city trends in PM concentrations.
Pope et al. 2004a	Cohort study examining ambient PM exposure and cardiovascular mortality in 319,000-500,000 persons in the American Cancer Society study, with 16 years follow-up across U.S. urban areas.	Cardiovascular mortality: ischemic heart disease, dysrhythmias, heart failure, and cardiac arrest.	A 10 µg/m <sup>3</sup> increase in PM <sub>2.5</sub> was associated with 8-18% increases in mortality due to ischemic heart disease, dysrhythmias, heart failure, and cardiac arrest.	Not assessed
Abbey et al. 1999	Cohort study examining ambient PM <sub>10</sub> exposure, total suspended sulfates, SO <sub>2</sub> , O <sub>3</sub> , and NO <sub>2</sub> to mortality in 6,338 nonsmoking California Seventh Day Adventists with 19 yr follow-up.	Cardiopulmonary mortality	No associations.	No associations.
Hoek et al. 2002	Cohort study examining ambient traffic related air pollutant exposure (black smoke and NO <sub>2</sub> ) and cause specific mortality in 5000	Cardiopulmonary mortality	Cardiopulmonary mortality was associated with living near high traffic density (100 m to freeway or 50 m to major urban road) adjusted RR= 1.95 (95% CI: 1.09, 3.52), and was	Cardiopulmonary mortality was similarly associated with an increase of 30 µg/m <sup>3</sup> background plus

Studies	Design and Population	Outcomes	Findings for PM Mass and Components	Findings for Gases
	persons with 8 years follow-up in the Netherlands Cohort Study on Diet and Cancer.		associated with an increase of 10 $\mu\text{g}/\text{m}^3$ black smoke from background (central sites) + local sources (street proximity), RR 1.71 (95% CI: 1.10, 2.67).	local $\text{NO}_2$ , RR 1.81 (95% CI: 0.98, 3.34)
Kunzli et al. 2004	Cross-sectional study on the relationship between ambient $\text{PM}_{2.5}$ and carotid artery intima-media thickness (CIMT) using baseline data from two clinical trials in Los Angeles. Annual mean $\text{PM}_{2.5}$ exposure was estimated using data from 23 monitoring stations linked to home addresses with geostatistical models.	Carotid artery intima-media thickness	For each increase of annual mean 10 $\mu\text{g}/\text{m}^3$ $\text{PM}_{2.5}$ , CIMT increased by 5.9% (95% CI: 1, 11%). Adjustment for age reduced the coefficients, but further adjustment for covariates indicated robust estimates in the range of 3.9 to 4.3%.	Estimates for $\text{O}_3$ linked to zip code centroids were positive in relation to CIMT but not significant and smaller than $\text{PM}_{2.5}$ .
<b>Cardiac Ischemia and Related Outcomes</b>				
Pekkanen et al. 2002	Panel study examining ambient PM, $\text{NO}_2$ , CO exposure and ischemia during 342 submaximal exercise tests in 45 subjects with CHD in Helsinki, Finland.	Electrocardiographic ST segment depression over 0.1 mV	Increased risk for ST depression (72 events) was associated with a change of lag 2 1000 particles/ $\text{cm}^3$ of accumulation mode PM ( $\text{NC}_{0.1-1}$ ) OR=3.29 (95% CI: 1.57, 6.92), and 10,000 particles/ $\text{cm}^3$ of UFP ( $\text{NC}_{0.01-0.1}$ ) OR=3.14 (95% CI: 1.56, 6.32). UFP was independent of associations with a 10 $\mu\text{g}/\text{m}^3$ increase in lag 2 $\text{PM}_{2.5}$ [OR=2.84, 95% CI: 1.42, 5.66].]	$\text{NO}_2$ and CO were also associated with an increased risk for ST depression.

Studies	Design and Population	Outcomes	Findings for PM Mass and Components	Findings for Gases
de Hartog et al, 2003	Panel study examining ambient exposure to PM and NO <sub>2</sub> , SO <sub>2</sub> and CO on HRV and BP in 131 subjects with CHD in Helsinki, Finland, Amsterdam, the Netherlands, Erfurt, Germany.	Cardiorespiratory symptoms: chest pain, shortness of breath, avoidance of activities	A 10 µg/m <sup>3</sup> increase in PM <sub>2.5</sub> associated with shortness of breath and avoidance of activities.	Not assessed
Peters et al. 2004	Case-crossover study examining ambient traffic related air pollution exposure and myocardial infarction in 691 subjects from the Augsburg Myocardial Infarction Registry who had survived 24 h post infarct. Time-activity diary data on activities during the four days before symptom onset was used to assess traffic exposures.	Myocardial Infarction	Exposure to traffic was associated with onset of myocardial infarction one hour afterward, OR =2.92 (95% CI: 2.22, 3.83). A significant association was also seen for exposure to traffic two hours before onset, and there was evidence for effects up to six hours. Key exposures influencing overall associations with traffic included times spent in cars and in public transportation. Associations changed minimally adjusting for exercise, and there was no confounding by reports of extreme anger or joy.	As with PM, gases were not directly assessed, but traffic exposures involve pollutant gases as well as particles.
<b>Blood Pressure (BP)</b>				
Linn et al. 1999	Panel study in Los Angeles, California examining BP and lung function in 30	BP	Systolic BP increased 0.172 mm Hg for every 1 µg/m <sup>3</sup> increase in ambient lag 1 PM <sub>10</sub> ( <i>p</i> = 0.006). Diastolic BP	No association of BP with exposure to central site O <sub>3</sub> , NO <sub>2</sub> , or CO.

Studies	Design and Population	Outcomes	Findings for PM Mass and Components	Findings for Gases
	subjects with COPD, with only four consecutive days of air sampling: personal exposure to PM <sub>2.5</sub> , indoor and outdoor home PM <sub>2.5</sub> and PM <sub>10</sub> , and ambient PM <sub>10</sub> , O <sub>3</sub> , NO <sub>2</sub> , and CO.		increased 0.095 mm Hg for every 1 µg/m <sup>3</sup> increase in PM <sub>10</sub> (P = 0.03). Outdoor home PM <sub>10</sub> was similarly associated with BP, but no significant associations were reported for PM <sub>2.5</sub> or any indoor or personal PM measurement.	
Brauer et al. 2001	Panel study examining personal exposure over seven nonconsecutive days to PM <sub>2.5</sub> and sulfate, and ambient exposure to ambient PM <sub>2.5</sub> , PM <sub>10</sub> , sulfate, and gaseous pollutants, in relation to BP, HRV, and lung function in 16 COPD patients in Vancouver, Canada.	BP, HRV, supraventricular ectopic heartbeats (SVE)	Weak associations were observed between particle concentrations and increased SVE and with decreased systolic BP. Ambient PM <sub>10</sub> had the largest effect on cardiovascular endpoints and the only statistically significant association (SVE). Use of personal exposure measurements did not show a larger or more consistent effect.	CO was inversely associated with systolic BP and reduced estimates for ambient PM.
Ibald-Mulli et al. 2001	Retrospective analysis examining the relationship between ambient air pollution exposure (TSP, SO <sub>2</sub> and CO) and blood pressure in 2607 men and women ages 25-64 y from a general population survey in Augsburg, Germany.	Systolic BP	A 90 µg/m <sup>3</sup> increase in TSP was associated with an increase in systolic BP of 1.79 mm Hg (95% CI: 0.63, 2.95). In subgroups with high plasma viscosity levels or increased heart rates, systolic BP increased by 6.93 mm Hg (95% CI: 4.31, 9.75) and 7.76 mm Hg (95% CI: 5.70, 9.82) in association with TSP, respectively	An 80 µg/m <sup>3</sup> increase in SO <sub>2</sub> was associated with an increase in systolic BP of 0.74 mm Hg (95% CI: 0.08, 1.40).
Ibald-Mulli et al. 2004	Panel study examining ambient exposure to PM and NO <sub>2</sub> , SO <sub>2</sub> and CO on HRV	BP and HR (Ibald-Mulli et al. 2004);	A small decrease in systolic BP (−0.72 mm Hg, 95% CI: −1.92, 0.49) and diastolic BP (−0.70 mm Hg, 95% CI:	The magnitude and significance of inverse BP associations with CO were

Studies	Design and Population	Outcomes	Findings for PM Mass and Components	Findings for Gases
	and BP in 131 subjects with CHD in Helsinki, Finland, Amsterdam, the Netherlands, Erfurt, Germany.		–0.02, –1.38) was found to be associated with a 5-day average increase of 10,000 ultrafine particles/cm <sup>3</sup> (PM <sub>0.01-0.1</sub> ). Slightly stronger and more significant associations were found for accumulation mode particle number concentration (PM <sub>0.1-1.0</sub> ), but smaller associations were found for a 10 µg/m <sup>3</sup> increase in PM <sub>2.5</sub> mass. Small decreases in HR were also found for PM exposures.	similar to PM <sub>0.1-1.0</sub> . A small decrease in HR (–0.40 beats/min, 95% CI: –0.82, 0.01) was found for an increase of lag 1 5 µg/m <sup>3</sup> SO <sub>2</sub> .
Zanobetti et al. 2004	Panel study examining the association between ambient PM <sub>2.5</sub> and blood pressure among 62 patients with pre-existing heart disease using data from 631 repeated visits for cardiac rehabilitation in Boston.	BP	Increasing from the 10 <sup>th</sup> to the 90 <sup>th</sup> percentile in 5-day mean PM <sub>2.5</sub> resulted in a 2.8 mm Hg (95% CI: 0.1, 5.5) increase in systolic, a 2.7 mm Hg (95% CI, 1.2 to 4.3) increase in diastolic, and a 2.7-mm Hg (95% CI: 1.0, 4.5) increase in mean arterial blood pressure.	Not assessed.
<b>Autonomic Control of Cardiac Rhythm</b>				
Holguin et al. 2003	Panel study in Mexico City examining indoor and outdoor nursing home measurements of PM <sub>2.5</sub> , and ambient exposure to O <sub>3</sub> , NO <sub>2</sub> , CO, SO <sub>2</sub> in relation to HRV in 34 elderly residents followed	HRV, frequency domain	A 10 µg/m <sup>3</sup> increase in predicted personal PM <sub>2.5</sub> was associated with a 5.0% decrease in high frequency HRV (β = –0.049, 95% CI: –0.090, –0.007). Associations with indoor PM <sub>2.5</sub> were stronger than outdoor home PM <sub>2.5</sub> . Among 13 subjects with hypertension, the association with predicted personal	O <sub>3</sub> was inversely associated with high and low frequency HRV among 13 subjects with hypertension (2% decrease per 10 ppb O <sub>3</sub> ), but this association was confounded by PM <sub>2.5</sub> .

Studies	Design and Population	Outcomes	Findings for PM Mass and Components	Findings for Gases
	every other day for 3 months. Personal PM <sub>2.5</sub> was predicted using indoor and outdoor home PM <sub>2.5</sub> plus time-activity data		PM <sub>2.5</sub> was stronger (−7.1%).	
Pope et al. 2004b	Panel study of ambient exposure to PM and HRV and blood markers in 88 elderly subjects living in Salt Lake, and Provo/Orem Utah	HRV and blood markers of inflammation (see below)	A 100 µg/m <sup>3</sup> increase in PM <sub>2.5</sub> was associated with a 35 (SE = 8) msec decrease in SDNN, and a 42 (SE = 11) msec decrease in r-MSSD	Not assessed
Magari et al. 2001; 2002a; 2002b	Panel study examining personal exposure to PM in relation to HRV in 20 (Magari et al. 2002a) 40 (Magari et al. 2001) and 39 (Magari et al. 2002b) healthy boilermakers exposed to welding fumes and residual oil fly ash.	HRV	Each 100 µg/m <sup>3</sup> increase in 3-hr average PM <sub>2.5</sub> (laser photometer light scatter) was associated with a 1.4% (95% CI: −2.1, −0.6%) decrease in 5-min SDNN in the 20 subjects (Magari et al. 2002a). In the 40 subjects, each 1 mg/m <sup>3</sup> increase in 4-hr average PM <sub>2.5</sub> was associated with a 2.66% (95% CI: −3.75, −1.58%) decrease in 5-min SDNN (Magari et al. 2001). However, in 39 of the same subjects, PM <sub>2.5</sub> metals on filters, lead and vanadium, were associated with an increase in workday average of the 5-min SDNN (Magari et al. 2002b).	Not assessed
Riediker et al. 2004	Panel study of in-vehicle exposure to PM and HRV and blood markers of inflammation in 9 healthy	HRV and blood markers of inflammation (see below)	In-vehicle 10 µg/m <sup>3</sup> PM <sub>2.5</sub> increase was associated with increased ectopic beats throughout exposure (20%, <i>p</i> = 0.005). PM <sub>2.5</sub> was positively associated with	NO <sub>2</sub> and CO were not significant



Studies	Design and Population	Outcomes	Findings for PM Mass and Components	Findings for Gases
	male North Carolina Highway Patrol troopers.		heart beat cycle length (6%, $p = 0.01$ ) as well as HF HRV and SDNN the next morning following exposure.	
Chan et al. 2004	Panel study in Taipei, Taiwan examining personal exposure to submicrometer particles and HRV over one 16-hr daytime period in 9 young healthy adults ages 19-29 y (2 females) and 10 older male subjects ages 42-97 y with lung function impairments ( $FEV_1/FVC < 85\%$ ).	HRV	Personal exposure to $NC_{0.02-1}$ was associated with decreased in both time-domain and frequency-domain HRV indices. In young subjects, a 10,000 particles/cm <sup>3</sup> increase in the last 1-4 hr average $NC_{0.02-1}$ was associated with 0.68-1.35% decrease in SDNN, 1.85-2.58% decrease in r-MSSD. In the older panel they found 10,000 particles/cm <sup>3</sup> increase in the last 1-3 hr average $NC_{0.02-1}$ was associated 1.72-3.00% decreases in SDNN, and 2.72-4.65% decreases in r-MSSD. There were similar associations for high and low frequency-domain indices.	Not assessed
Tarkiainen et al. 2003	Panel study in Kuopio, Finland, examining personal exposure to carbon monoxide and HRV in 6 subjects with CHD followed for three separate 24-hr ambulatory monitoring periods.	HRV	Not assessed	r-MSSD increased by 2.4 ms ( $p = 0.03$ ) with exposure to CO ( $> 2.7$ ppm).
Peters et al. 2000	Panel study of arrhythmias in 100 subjects in eastern Massachusetts with implanted defibrillators	Defibrillator discharge interventions for ventricular	Only 6 subjects with $\geq 10$ defibrillator discharges had increased arrhythmias associated black carbon and $PM_{2.5}$ , which showed a weaker association.	26-ppb increase in $NO_2$ lagged 1 d was associated with increased defibrillator interventions in the full

Studies	Design and Population	Outcomes	Findings for PM Mass and Components	Findings for Gases
	(63,628 person-days of follow-up) with ambient measurements of PM mass, black carbon, NO <sub>2</sub> , CO, O <sub>3</sub> and SO <sub>2</sub>	tachycardias or fibrillation (33 subjects with at least one).	Both PM metrics were confounded by NO <sub>2</sub> , but the effect estimate of NO <sub>2</sub> was unchanged.	panel (OR 1.8; 95% CI: 1.1, 2.9). Subjects with ≥10 defibrillator discharges had increased arrhythmias associated with CO and NO <sub>2</sub> . across several lags
<b>Systemic Inflammation and Thrombosis</b>				
Seaton et al. 1999	Panel study examining 3 day* personal exposure and city center ambient exposure to PM <sub>10</sub> on hematological factors in 112 elderly subjects in Belfast and Edinburgh, UK.  *(personal exposure estimated from a one 24 hour personal exposure measurement)	Hematological factors: hemoglobin, packed red cells, red blood cell count, platelets, white blood cell count, CRP, fibrinogen, factor VII, interleukin-6	An increase of 100 µ/m <sup>3</sup> in personal PM <sub>10</sub> and ambient PM <sub>10</sub> exposure resulted in significant decreased mean percentage changes of ≤ 1% in hemoglobin concentration, packed cell volume, and red blood cell count. Only personal PM <sub>10</sub> was associated with an 11% decrease in platelets and a 7% decrease in Factor VII. CRP increased with ambient PM <sub>10</sub> (+147%, 95% CI: 20, 477), but not with personal PM ( <i>p</i> = 0.73). Fibrinogen decreased with ambient PM <sub>10</sub> (-9%, 95% CI: -19, 0).	Not assessed
Schwartz et al. 2001	Cross-sectional study examining the relationship between ambient PM <sub>10</sub> , NO <sub>2</sub> , SO <sub>2</sub> , and blood biomarkers using data from a cohort study (NHANES III).	Fibrinogen, and platelet and white blood cell counts	For an interquartile range change in PM <sub>10</sub> (26 µg/m <sup>3</sup> ) the relative odds for being above the 90th percentile of fibrinogen was 1.77 ( 95% CI: 1.26, 2.49), platelets, 1.27 ( 95% CI: 0.97, 1.67) and white blood cells, 1.64 ( 95% CI: 1.17, 2.30).	SO <sub>2</sub> was significantly associated with white cell counts, NO <sub>2</sub> with platelet counts and fibrinogen, but both gases were confounded by PM <sub>10</sub> .
Pekkanen et al.	Cross-sectional study examining the association between ambient PM <sub>10</sub> ,	Fibrinogen	No association between PM <sub>10</sub> and fibrinogen was seen after adjustment for confounders.	NO <sub>2</sub> increase from the 10 <sup>th</sup> to the 90 <sup>th</sup> percentile was associated with a 1.5%

Studies	Design and Population	Outcomes	Findings for PM Mass and Components	Findings for Gases
2000	NO <sub>2</sub> , CO, SO <sub>2</sub> , O <sub>3</sub> and fibrinogen among 7205 subjects in London at baseline enrollment in a cohort study.			higher fibrinogen concentration (95% CI: 0.4% to 2.5%). Similar increase for CO resulted in 1.5% higher fibrinogen concentration (95% CI: 0.5%, 2.5%). No association with SO <sub>2</sub> or O <sub>3</sub> .
Peters et al. 2001b Peters et al. 1997a	Cohort study in Augsburg, Germany, examining relationships of ambient TSP, SO <sub>2</sub> , and CO exposure to CRP in 631 men age 45-64 with no history of MI at their baseline assessment. There were two CRP measurements three years apart.	CRP	An increase of 26 µg/m <sup>3</sup> (5-day mean) in TSP increased the odds of observing a CRP level above the 80 <sup>th</sup> percentile, OR = 1.31 (95% CI: 1.09, 1.56). CRP and plasma viscosity (Peters et al. 1997a) were increased during an air pollution episode in 1985.	An increase of 30 µg/m <sup>3</sup> (5 day mean) in SO <sub>2</sub> increased the odds of observing a CRP level above the 90 <sup>th</sup> percentile, OR = 1.24 (95% CI: 1.03, 1.49).
Pope et al. 2004b	Panel study of ambient exposure to PM and HRV and blood markers in 88 elderly subjects living in Salt Lake, and Provo/Orem Utah.	HRV (see above) and CRP, white blood cell count, whole blood viscosity, granulocytes, lymphocytes, monocytes, basophils, eosinophils, red blood cells, platelets	A 100 µg/m <sup>3</sup> increase in PM <sub>2.5</sub> was associated with a 0.81 (SE = 0.17) mg/dl increase in CRP. One subject's data had a strong influence on estimates. There was no association with other outcomes.	Not assessed

Studies	Design and Population	Outcomes	Findings for PM Mass and Components	Findings for Gases
Riediker M et al. 2004	Panel study of in vehicle exposure to PM and HRV and blood markers of inflammation in 9 healthy male North Carolina Highway Patrol troopers.	HRV (see above) and CRP, plasminogen, von Willebrand factor, Lymphocyte count, lymphocytes, neutrophils, hematocrit, red blood cell indices, uric acid	In-vehicle 10 $\mu\text{g}/\text{m}^3$ PM <sub>2.5</sub> increase was associated with decreased lymphocytes ( $-11\%$ , $p = 0.03$ ), increased red blood cell indices ( $1\%$ , $p = 0.03$ ), neutrophils ( $6\%$ , $p = 0.04$ ), CRP ( $32\%$ , $p = 0.02$ ), and von Willebrand factor ( $12\%$ , $p = 0.02$ ).	NO <sub>2</sub> and CO were not significant

<sup>a</sup> The focus is on cardiovascular outcomes. Although some studies may have examined other outcomes, they are not reported.

## **FIGURE LEGENDS**

Figure 1. Hypothesized pathways leading to adverse cardiovascular health effects from exposure to ultrafine particles

Figure 1.

